

## A Study on the Estimator Distribution for the Expected Value of a Compound Periodic Poisson Process with Power Function Trend

Nurul Indah Safitri\*, I Wayan Mangku and Hadi Sumarno  
Department of Mathematics, IPB University, Bogor 16680, Indonesia  
Email: pasca\_nurul@apps.ipb.ac.id

### Abstract

This article discusses the estimation for the expected value, also called the mean function, of a compound periodic Poisson process with a power function trend. The aims of our study are, first, to modify the existing estimator to produce a new estimator that is normally distributed, and, second, to determine the smallest observation interval size such that our proposed estimator is still normally distributed. Basically, we formulate the estimator using the moment method. We use Monte Carlo simulation to check the distribution of our new estimator. The result shows that a new estimator for the expected value of a compound periodic Poisson process with a power function trend is normally distributed and the simulation result shows that the distribution of the new estimator is already normally distributed at the length of 100 observation interval for a period of 1 unit. This interval is the smallest size of the observation interval. The Anderson-Darling test shows that when the period is getting larger, the  $p$ -value is also getting bigger. Therefore, the larger period requires a wider observation interval to ensure that the estimator still has a normal distribution.

**Keywords:** moment method; normal distribution; Poisson process; the smallest observation interval.

### Abstrak

*Pada artikel ini dibahas tentang pendugaan fungsi nilai harapan Proses Poisson periodik majemuk dengan tren fungsi pangkat. Tujuan penelitian kami adalah, pertama, memodifikasi penduga yang telah ada untuk menghasilkan penduga baru yang memiliki distribusi normal. Kedua, menentukan ukuran interval pengamatan terkecil sehingga penduga yang diusulkan masih berdistribusi normal. Pada dasarnya, penduga yang kami usulkan diformulasi menggunakan metode momen. Kami menggunakan metode simulasi Monte Carlo untuk memeriksa sebaran distribusinya. Hasil menunjukkan bahwa penduga yang baru untuk fungsi nilai harapan Proses Poisson periodik majemuk dengan tren fungsi pangkat memiliki distribusi normal. Hasil simulasi menunjukkan bahwa penduga baru telah berdistribusi normal pada panjang interval pengamatan 100 untuk periode sebesar 1 satuan. Interval pengamatan ini merupakan ukuran interval pengamatan terkecil. Selain itu, hasil uji Anderson-Darling menunjukkan bahwa ketika periode semakin besar maka  $p$ -value juga semakin besar. Oleh karena itu, periode yang lebih besar memerlukan interval pengamatan yang lebih panjang untuk menjamin penduga yang kami usulkan tetap berdistribusi normal.*

**Kata Kunci:** metode momen; distribusi normal; proses Poisson; interval pengamatan terkecil.

2020MSC: 62E17

## 1. INTRODUCTION

We consider the problem of estimating the expected value, also called the mean function, of a compound Poisson process when the intensity function of the Poisson component is a periodic or cyclic function plus a power function trend. Since the periodic compound Poisson process has wide applications in applied sciences, it is needed to know the distribution of this process. However, it is

---

\* Corresponding author

Submitted March 7<sup>th</sup>, 2022, Revised June 30<sup>th</sup>, 2022, Accepted for publication July 20<sup>th</sup>, 2022.

©2022 The Author(s). This is an open-access article under CC-BY-SA license (<https://creativecommons.org/licenses/by-sa/4.0/>)

hard to estimate the exact distribution of this process. Hence, some researchers try to estimate two aspects of this process, namely its mean function ( [1] [2] [3] [4] [5] [6]) and its variance function ( [3] [6] [7] [8]).


Erliana et al. [11], Maulidi, et al. [12] and Mulidi et al. [13] discussed the power function trend of a non-homogeneous Poisson process. Mangku [14], Maulidi et al. [15], and Abdullah et al. [16] estimated the intensity of a cyclic Poisson process with the power function of a non-homogeneous Poisson process. The confidence interval for the expected value function had been estimated in [17] and [18]. The Poisson process had been applied to ore flow in a belt conveyor system [19], in the probability distribution of COVID-19 [20], and in pricing catastrophe bonds [21].

Formulation of consistent estimation of the mean (expected value) function of purely cyclic compound Poisson process was first done in [1]. In [2], this estimator was slightly modified in order to obtain asymptotic bias and asymptotic variance of the estimator. In [4], a consistent estimator for the mean (expected value) function of a compound periodic Poisson process with power function trend was formulated. The asymptotic bias and asymptotic variance of this estimator also have been formulated in [4], [9], and [10].

This research is a continuation of Sari et al. [4]. Sari et al. [4] do not assume any parametric form for the estimator for the mean function whereas in this research the estimator has a distribution. We will formulate a new estimation for the expected value of a compound periodic Poisson process with power function trend with the objectives as follows. (i) To modify the estimator formulated in [4] to obtain a new estimator which is normally distributed. (ii) To find the smallest size of the observation interval of the Poisson process such that the estimator is still normally distributed.

## 2. METHODS

Figure 1 shows our research framework. We formulate the new estimator using the moment method and use Monte Carlo simulation to check the distribution of the new estimator.



**Figure 1.** The research framework.

The steps in this research include: (1) preliminary study and (2) the main study. The preliminary study comprises studying the compound periodic Poisson process with power function trend and exploring mathematical foundations to develop a new estimator from the previous theories. The main study involves modifying the existing estimator and investigating the normality of the new estimator; as well as obtaining the smallest observation interval size of the process where the new estimator is still normally distributed.

### 3. RESULTS AND DISCUSSION

In this section, formulation for the expected value (mean) function of a compound cyclic Poisson process with power function trend is discussed.

#### 3.1 Formulation of the Mean Function

Let  $\{N(t), t \geq 0\}$  be a nonhomogeneous Poisson process having (unknown) intensity function  $\lambda$ . It is assumed that intensity function  $\lambda$  has two components, namely a cyclic (periodic) component,  $\lambda_c$ , with (known) period  $\tau > 0$  and a power function trend component. In other words, for all  $s \geq 0$ , the intensity function  $\lambda$  can be written as ([4])

$$\lambda(s) = \lambda_c(s) + as^b, \tag{1}$$

with  $\lambda_c$  is the cyclic component with period  $\tau$  and  $as^b$  is the power function trend, with  $a > 0$  is the slope of the power function trend. It is assumed that  $b$  is a known real number and  $0 < b < \frac{1}{2}$ . It is not assumed any parametric form for the cyclic component  $\lambda_c$ , except that it is cyclic or periodic, which satisfies

$$\lambda_c(s) = \lambda_c(s + k\tau) \tag{2}$$

for all  $s \geq 0$  and  $k \in \mathbb{N}$ , where  $\mathbb{N}$  denotes the set of natural numbers.

Let  $\{Y(t), t \geq 0\}$  be a process where

$$Y(t) = \sum_{i=1}^{N(t)} X_i \tag{3}$$

with  $\{X_i, i \geq 1\}$  is a sequence of i.i.d. (independent and identically distributed) random variables having mean  $\mu < \infty$  and variance  $\sigma^2 < \infty$ , and also independent of the process  $\{N(t), t \geq 0\}$ . The Process  $\{Y(t), t \geq 0\}$  is called a compound cyclic Poisson process with power function trend [4].

Let for some  $\omega \in \Omega$ , a single realization  $N(\omega)$  of the process  $\{N(t), t \geq 0\}$  is observed on an interval  $[0, m]$ ,  $m$  is a real number. Furthermore, suppose that for each data point in the realization  $N(\omega) \cap [0, m]$ , say  $i$ -th data point,  $i = 1, 2, \dots, N[0, m]$ , its corresponding random variable  $X_i$  is also observed.

Let  $\psi(t)$  denotes the expected value (mean) function of the considered process.  $\psi(t)$  can be written as

$$\psi(t) = E[Y(t)] = E[N(t)]E[X_i] = \Lambda(t)\mu, \tag{4}$$

with  $\mu = E(X_i)$  and

$$\Lambda(t) = \int_0^t \lambda(s) ds. \tag{5}$$

Let  $t_r = t - \left\lfloor \frac{t}{\tau} \right\rfloor \tau$  where  $\lfloor x \rfloor$  denotes the biggest integer that is less than or equal to  $x \in \mathbb{N}$ , and

$k_{t,\tau} = \left\lfloor \frac{t}{\tau} \right\rfloor$ . Then for any real number  $t \geq 0$ , we can write  $t = k_{t,\tau}\tau + t_r$  with  $0 \leq t_r < \tau$ . Let  $\tau\theta = \int_0^\tau \lambda_c(s) ds$

, which can be written as  $\Lambda_c(t_r) + \Lambda_c^c(t_r)$  with  $\Lambda_c(t_r) = \int_0^{t_r} \lambda(s) ds$  and  $\Lambda_c^c(t_r) = \int_{t_r}^\tau \lambda(s) ds$ . It is

assumed that  $\theta > 0$ . By using the above notations and substituting (1) into (5), for any  $t \geq 0$ ,  $\Lambda(t)$  can be formulated as

$$\Lambda(t) = (k_{t,\tau} + 1)\Lambda_c(t_r) + k_{t,\tau}\Lambda_c^c(t_r) + \frac{a}{b+1}t^{b+1}. \tag{6}$$

By substituting (6) into the r.h.s. of (4), we have

$$\psi(t) = \left( (k_{t,\tau} + 1)\Lambda_c(t_r) + k_{t,\tau}\Lambda_c^c(t_r) + \frac{a}{b+1}t^{b+1} \right) \mu. \tag{7}$$

### 3.2 Formulation of the Estimator for the Mean Function

After formulating the mean function, we formulate the estimator for the expected value (mean) function of the process as follows. Let  $n^{1+\delta} = m$ , where  $\delta$  be an arbitrarily small positive real number. For some reason, to estimate  $a$  we use the realization of the Poisson process observed in  $[0, m]$ , while to estimate the other quantities we use the realization of the Poisson process observed in  $[0, n]$ . The estimator of the mean function can be written as

$$\hat{\psi}_{n,b}(t) = \left( (k_{t,\tau} + 1)\hat{\Lambda}_{c,n,b}(t_r) + k_{t,\tau}\hat{\Lambda}_{c,n,b}^c(t_r) + \frac{\hat{a}_{m,b}}{b+1}t^{b+1} \right) \hat{\mu}_n, \tag{8}$$

where

$$\hat{\Lambda}_{c,n,b}(t_r) = \frac{(1-b)\tau^{1-b}}{n^{1-b}} \sum_{k=1}^{k_{n,\tau}} \frac{1}{k^b} N([k\tau, k\tau + t_r]) - \hat{a}_{m,b}(1-b)n^b t_r,$$

$$\hat{\Lambda}_{c,n,b}^c(t_r) = \frac{(1-b)\tau^{1-b}}{n^{1-b}} \sum_{k=1}^{k_{n,\tau}} \frac{1}{k^b} N([k\tau + t_r, k\tau + \tau]) - \hat{a}_{m,b}(1-b)n^b(\tau - t_r),$$

$$\hat{a}_{m,b} = \frac{(1+b)N([0,m])}{m^{1+b}} - \frac{(1+b)}{m^b} \tilde{\theta}_n,$$

$$\tilde{\theta}_n = \frac{(1-b)}{n^{1-b} \tau^b b^2} \sum_{k=1}^{k_{n,\tau}} \frac{1}{k^b} N([k\tau, k\tau + \tau]) - \frac{(1-b)(1+b)n^b N([0,n])}{n^{1+b} b^2}, \text{ and}$$

$$\hat{\mu}_n = \frac{1}{N[0,n]} \sum_{i=1}^{N[0,n]} X_i, \text{ if } N([0,n]) > 0 \text{ and } \hat{\mu}_n = 0 \text{ if } N([0,n]) = 0.$$

The estimator for the global intensity function  $\theta$  is a modification of the estimator formulated in [4]. In this paper, the normality of the estimator given in (8) was investigated using Monte Carlo simulation. We also determine the smallest size of the observation interval such that the distribution of the estimator is already normal.

### 3.3 The Monte Carlo Simulation

The simulation was carried out with the help of R software. The simulation is aimed at checking normality distribution and determining the smallest  $n$  value where the distribution of the estimator is close to the normal distribution. The programming stage begins with generating the realization of a compound periodic Poisson process with a power function trend with the formulation of the intensity function as follows.

$$\lambda(t) = \sin\left(\frac{2\pi t}{\tau}\right) + 1 + 0.07t^{2/5}. \tag{10}$$

In this study, the normality test was carried out using a QQ-plot graph analysis approach. The data is normally distributed if the spread of each point approaches the diagonal line, hence the assumption of normality is met. The simulation test results can be seen in Figures 2-5.






Figure 2. Cont.




(c)

**Figure 2.** Normality of  $\hat{\Psi}_{n,b}(t)$  for  $\tau = 1$ . (a) Observation interval at  $[0, 25]$ ; (b) Observation interval at  $[0, 75]$ ; (c) Observation interval at  $[0, 100]$ .




(a)




(b)

**Figure 3.** Normality of  $\hat{\Psi}_{n,b}(t)$  for  $\tau = 5$ . (a) Observation interval at  $[0, 100]$ ; (b) Observation interval at  $[0, 200]$ .




(a)



(b)

**Figure 4.** Normality of  $\hat{\Psi}_{n,b}(t)$  for  $\tau = 10$ . (a) Observation interval at  $[0, 300]$ ; (b) Observation interval at  $[0, 500]$ .



**Figure 5.** Normality of  $\hat{\Psi}_{n,b}(t)$  for  $\tau = 30$ . (a) Observation interval at  $[0, 900]$ ; (b) Observation interval at  $[0, 1200]$ .

Based on the simulation results, in the observation intervals  $[0, 25]$ ,  $\tau = 1$  (Figure 2(a)) and  $[0, 75]$ ,  $\tau = 1$  (Figure 2(b)), the spread of each estimator point is still visible away from the diagonal line. Meanwhile, in the observation interval  $[0, 100]$ ,  $\tau = 1$  (Figure 2(c)), the spread of each estimator point already approaches the diagonal line. This illustration shows that the estimator for the mean function of a compound periodic Poisson process with a power function trend is getting closer to the normal distribution for a longer interval of observation. Furthermore, the Anderson-Darling test was carried out to check the normality of the estimator based on the  $p$ -value. The  $p$ -value of the normality simulation results of the estimator for the mean function is shown in Table 1.

Table 1 shows that the  $p$ -value when the period of 1 unit for the observation intervals  $[0, 25]$  and  $[0, 75]$  is less than 0.05. It means that the estimator of the mean function is not normally distributed. Meanwhile, for the same period of 1 unit with an observation interval at  $[0, 100]$ , the  $p$ -value is more than 0.05. That is, the estimator of the mean function begins to spread normally. So,  $[0, 100]$  is the smallest length of the observation interval where the distribution of the estimator for the mean function is normally distributed. Furthermore, note that the  $p$ -value at the length of the observation interval  $[0, 100]$  when the period of 1 unit is greater than the  $p$ -value at the length of the observation interval  $[0, 100]$  when the period is 5. This is because when the period is getting bigger, the error generated is also getting bigger. When the period is getting longer, then the  $p$ -value is also getting bigger. This means that a longer period requires a wider observation interval to ensure that the estimator is still normally distributed.

**Table 1.** The simulation of normality for the estimator of mean function

Period ( $\tau$ )	Observation Interval $[0, n]$	$p$ -Value
1	$[0,25]$	$<0.05$
	$[0,75]$	$<0.05$
	$[0,100]$	0.1231
5	$[0,100]$	0.0617
	$[0,200]$	0.0955
10	$[0,300]$	0.2994
	$[0,500]$	0.3564
30	$[0,900]$	0.4582
	$[0,1200]$	0.9608

#### 4. CONCLUSIONS

Our study proposes a new estimator for the expected value (mean) function of a compound periodic Poisson process with a power function trend that has a normal distribution. The simulation result shows that our new estimator is already normally distributed at the length of 100 observation interval for a period of 1 unit. This is the smallest size of observation interval. The Anderson-Darling test shows that as the period gets larger, the  $p$ -value also gets larger. It means that, the larger the period, it requires wider observation interval to ensure that the estimator is still normally distributed.

#### REFERENCES

- [1] Ruhiyat, I. W. Mangku and , I. G. P. Purnaba, "Consistent estimation of the mean function of a compound cyclic Poisson process," *Far East Journal of Mathematical Science*, vol. 77, pp. 183-194, 2013.
- [2] I. W. Mangku, Ruhiyat and I. G. P. Purnaba, "Statistical properties of an estimator for the mean function of a compound cyclic Poisson process," *Far East Journal of Mathematical Science*, vol. 82, pp. 227-237, 2013.
- [3] I. W. Mangku, F. Sakinah and Ruhiyat, "Estimating the mean and variance function of a compound Poisson process having power function intensity," *Far East Journal of Mathematical Sciences*, vol. 100, pp. 1455-1465, 2016.
- [4] I. F. Sari, I. W. Mangku and H. Sumarno, "Estimating the mean function of a compound cyclic Poisson process in the presence of power function trend," *Far East Journal of Mathematical Sciences*, vol. 100, pp. 1825-1840, 2016.
- [5] B. A. Wibowo, I. W. Mangku and Siswadi, "Statistical properties of an estimator for the mean function of a compound cyclic Poisson process in the presence of linear trend," *Arab Journal of Mathematical Sciences*, vol. 23, pp. 173-185, 2017.
- [6] I. M. Y. E. Prasetya, I. W. Mangku and H. Sumarno, "Estimating the mean and variance of a compound Poisson process with the Poisson intensity obtained as exponential of the linear function," *Far East Journal of Mathematical Sciences*, vol. 102, pp. 721-729, 2017.
- [7] F. I. Makhmudah , I. W. Mangku and H. Sumarno, "Estimating the variance function of a compound cyclic Poisson process," *Far East Journal of Mathematical Sciences*, vol. 100, pp. 911-922, 2016.
- [8] S. Abdullah , I. W. Mangku and Siswadi, "Estimating the variance function of a compound cyclic Poisson process in the presence of linear trend," *Far East Journal of Mathematical Sciences*, vol. 102, pp. 559-572, 2017.
- [9] M. W. Utama, I. W. Mangku and B. P. Silalahi, "Asymptotic distribution of an estimator for variance function of a compound periodic Poisson process with power function trend," *Jurnal Teori dan Aplikasi Matematika*, vol. 6, no. 4, pp. 1081-1095, 2022. doi:<https://doi.org/10.31764/jtam.v6i4.10213>.
- [10] N. Valentika, I. W. Mangku and W. Erliana, "Strong consistency and asymptotic distribution of estimator for the intensity function having form of periodic function multiplied by power



- function trend of a Poisson process," *International Journal of Engineering and Management Research*, vol. 8, no. 2, pp. 232-236, 2018.
- [11] W. Erliana, I. W. Mangku and H. Sumarno, "Estimating the intensity obtained as the product of a periodic function with the power function trend of a non-homogeneous Poisson process," *Far East Journal of Mathematical Science*, vol. 90, pp. 163-172, 2014.
- [12] I. Maulidi, I. W. Mangku and H. Sumarno, "Strong consistency of a kernel-type estimator for the intensity obtained as the product of a periodic function with the power function trend of a non-homogeneous Poisson process," *British Journal of Applied Science and Technology*, Vols. 383-387, p. 9, 2015.
- [13] I. Maulidi, M. Ihsan and V. Apriliani, "The numerical simulation for asymptotic normality of the intensity obtained as a product of a periodic function with the power trend function of a nonhomogeneous Poisson process," *Desimal: Jurnal Matematika*, vol. 3, no. 271-278, p. 3, 2020.
- [14] I. W. Mangku, "Estimating the intensity of a cyclic Poisson process in the presence of additive and multiplicative trends," *Journal of Physics: Conference Series 893 (1)*, 012023, 2017.
- [15] I. Maulidi, W. A. B., N. Valentika, M. Syazali and V. Apriliani, "The confidence interval for the periodic intensity function in the presence of power function trend on the nonhomogeneous Poisson process," *CAUCHY: Jurnal Matematika Murni dan Aplikasi*, vol. 7, no. 1, pp. 78-83, 2021. doi: <http://dx.doi.org/10.18860/ca.v7i1.12848>.
- [16] S. Abdullah, S. Susilo, I. W. Mangku, F. Ikhsan, S. Ula and Y. Rukmayadi, "Algorithm for generating compound Poisson process which has nonhomogeneous Poisson process and exponential distribution components," in *The 1st International Multidisciplinary Conference on Education, Technology, and Engineering (IMCETE)*, 2020. doi: 10.2991/assehr.k.200303.059.
- [17] A. Fithry, I. W. Mangku and I. G. P. Purnaba, "Interval Kepercayaan untuk Fungsi Nilai Harapan dan Fungsi Ragam Proses Poisson Periodik Majemuk," *Jambura J. Math*, vol. 4, no. 1, p. 152-166, 2022. <https://doi.org/10.34312/jjom.v4i1.12180>.
- [18] I. Maulidi, B. A. Wibowo, N. Valentika, M. Syazali and V. Apriliani, "Confidence intervals for the mean function of a compound cyclic Poisson process in the presence of power function trend," *CAUCHY: Jurnal Matematika Murni dan Aplikasi*, vol. 7, no. 3, pp. 411-419, 2022.
- [19] P. Kruczek, M. Polak, A. Wylomanska, W. Kawalec and R. Zimroz, "Application of compound Poisson process for modelling of ore flow in a belt conveyor system with cyclic loading," *International Journal of Mining, Reclamation and Environment*, vol. 32, no. 6, pp. 376-391, 2018. <http://dx.doi.org/10.1080/17480930.2017.1388335>.
- [20] D. Munandar, S. Supian and Subiyanto, "Probability distributions of COVID-19 tweet posted trends uses a nonhomogeneous Poisson process," *International Journal of Quantitative Research and Modeling*, vol. 1, no. 4, pp. 119-238, 2020.
- [21] J. H. Sukono, R. A. Ibrahim, M. P. A. Saputra, Y. Hidayat and I. G. Prihanto, "Application of compound Poisson process in pricing catastrophe bonds: a systematic literature review," *MDPI Journal*, vol. 10, no. 15, p. 2668, 2022. doi:<https://doi.org/10.3390/math10152668>.