

# The Locating Chromatic Number for Amalgamation of Some Complete Graphs

Amanah Yulianti, Asmiati\*, Nur Hamzah, and Notiragayu  
Department of Mathematics, Faculty of Mathematics and Natural Sciences  
University Lampung, Bandar Lampung, Indonesia  
\*email: asmiati.1976@fmipa.unila.ac.id

## Abstract

The locating chromatic number of a graph is a combination of partition dimension and vertex coloring, where every two adjacent vertices are in different color classes, and all vertices have a unique color code. The amalgamation of  $a \geq 2$  complete graphs ( $K_n, n \geq 3$ ) denoted by  $aK_n$  is obtained by identifying one vertex from each complete graph. In this paper, we present a novel study, a topic that has not been extensively explored in previous research, on the locating chromatic number for the amalgamation of complete graph  $aK_n$  for  $2 \leq a \leq 6$  and  $n \geq 3$ .

**Keywords:** locating chromatic number; partition dimension; vertex coloring; color code; amalgamation of complete graph.

## Abstrak

Bilangan kromatik lokasi graf merupakan penggabungan dari dimensi partisi dan pewarnaan titik, yang mana setiap dua titik bertetangga berada dalam kelas warna yang berbeda dan semua titik mempunyai kode warna yang unik. Amalgamasi dari  $a \geq 2$  buah graf lengkap ( $K_n, n \geq 3$ ) dinotasikan dengan  $aK_n$  diperoleh dengan cara menyatukan satu titik dari setiap graf lengkap  $K_n$ . Pada paper ini didiskusikan hasil yang belum ada sebelumnya, yaitu bilangan kromatik lokasi amalgamasi graf lengkap  $aK_n$  untuk  $2 \leq a \leq 6$  dan  $n \geq 3$ .

**Kata Kunci:** bilangan kromatik lokasi, dimensi partisi, pewarnaan titik, kode warna, amalgamasi graf lengkap.

2020MSC: 05C12, 05C15

## 1. INTRODUCTION

The study of locating chromatic numbers is a fascinating subject in graph theory. This concept evolves from the ideas of partition dimensions and vertex coloring. The partition dimension of a graph is an extension of the metric dimension, which was first explored by Slater in 1975 [1]. The metric dimension of a graph is defined as the minimum number of members in the set representing a vertex's distances [2].

The locating chromatic number was introduced by Chartrand et al. in 2002 [3]. Let  $G = (V, E)$  be a connected graph, and  $q$  is a vertex coloring in  $G$  using the  $l$  colors of  $G$ , where  $q(u) \neq q(v)$  for any two adjacent vertices  $u, v$  in  $G$ . Suppose  $\Pi = \{Q_1, Q_2, \dots, Q_l\}$  is a partition of  $V(G)$  which is induced by coloring  $q$ . The color code  $q_\Pi(w)$  of  $w$  is  $l$ -pairs tuple  $(d(w, Q_1), d(w, Q_2), \dots, d(w, Q_l))$  with  $d(w, Q_i) = \min \{d(w, x) | x \in Q_i \text{ for } 1 \leq i \leq l$ . If all vertices in  $V(G)$  have different color codes, then  $q$  is called  $l$ -locating coloring of  $G$ . The locating chromatic number of a graph  $G$ , denoted by  $\chi_L(G)$  is the smallest  $l$  such that  $G$  has a  $l$ -locating coloring. Chartrand et al. [3] have obtained locating chromatic numbers from several graphs, including paths, cycles, and

\* Corresponding author

Submitted May 7<sup>th</sup>, 2024, Revised May 28<sup>th</sup>, 2024,

Accepted for publication May 29<sup>th</sup>, 2024, Published Online May 31<sup>st</sup>, 2024

©2024 The Author(s). This is an open-access article under CC-BY-SA license (<https://creativecommons.org/licence/by-sa/4.0/>)

double star graphs. One year later, Chartrand et al. [4] obtained the locating chromatic number of some graphs with locating chromatic number  $(n - 1)$ .

**Theorem 1.1.** (see[3]). *The locating chromatic number of a complete graph  $(K_n)$  is  $n$  for  $n \geq 2$ .*

Asmiati et al. [5] determined the locating chromatic number for the amalgamation of stars. In 2012, Asmiati found the locating chromatic number for firecracker graphs [6] and characterized graphs containing cycles with a locating chromatic number of three [7]. Baskoro and Asmiati [8] later characterized all trees with a locating chromatic number of three. Asmiati also obtained the locating chromatic number for nonhomogeneous caterpillars and firecracker graphs [9], certain barbell graphs [10], specific operations on origami graphs [11], barbell shadow path graphs [12], and the disjoint union of some double star graphs [13]. Irawan determined the locating chromatic number for subdivision firecracker graphs [14], various operations on generalized Petersen graphs [15][16], origami graphs [17] and provided a procedure for finding it [18], as well as the upper bounds of the locating chromatic numbers for shadow cycle graphs [19]. Behtoei found the locating chromatic number for Kneser graphs [20] and the Cartesian product of graphs [21]. Ghanem et al. [22] obtained the locating chromatic number for powers of paths and cycles. Furuya and Matsumoto [23] established upper bounds on the locating chromatic number for trees. Prawinasti et al. [24] determined the locating chromatic number for the split graph of the cycle. Damayanti et al. [25] found the locating chromatic number for some modified paths with cycles having a locating chromatic number of four. Finally, Rahmatalia et al. [26] determined the locating chromatic number for split graphs of paths.

Based on the literature study above, there is still no general formula for finding the locating chromatic number for any graph. So, we will discuss the locating chromatic number for amalgamation of complete graph and its barbell. The amalgamation of  $a \geq 2$  complete graphs  $(K_n, n \geq 3)$  denoted by  $aK_n$  is obtained by identifying one vertex from each complete graph. We call the identified vertex as the center (denoted by  $p$ ). Let  $V(aK_n) = \{ p, v_i^j \mid i = 1, 2, \dots, n - 1; j = 1, 2, 3, \dots, a; a \geq 2 \}$ . An example for amalgamation of complete graph  $4K_4$  with vertices label can be showed in Figure 1.




Figure 1. Amalgamation of complete graph  $4K_4$  with vertices label.

## 2. METHODS

To obtain the locating chromatic number of amalgamation of complete graph, we determine the lower and upper bound with construct the vertex coloring by looking at the graph structure. Vertex coloring starts with the smallest label so that a minimum vertex coloring is obtained that meets the location coloring requirements. If the lower bound and upper bound of the locating chromatic number of amalgamation of complete graph ( $aK_n$ ) are identical, for example  $x$ , then the locating chromatic number is  $\chi_L(aK_n) = x$ .

## 3. RESULTS AND DISCUSSION

**Theorem 3.1**  $\chi_L(aK_n) = n + 1$  for  $n \geq 3$  and  $a \in \{2,3\}$ .

**Proof.**

First, we determine the lower bound of the locating chromatic number of the graph  $aK_n$  for  $n \geq 3$  and  $a \in \{2,3\}$ . Amalgamation of complete graph  $aK_n$  contains  $K_n$ , by Theorem 1.1, we have  $\chi_L(aK_n) \geq n$ . Suppose that there is a  $n$ -locating coloring of  $aK_n$  for  $n \geq 3$  and  $a \in \{2,3\}$ . Since  $aK_n$  containing at least two  $K_n$ , then there are two vertices have the same color codes, a contrary. So,  $\chi_L(aK_n) \geq n + 1$ .

Next, we determined the upper bound of the locating chromatic number of  $aK_n$ . Let  $q$  be a  $(n + 1)$ -coloring in  $aK_n$  for  $n \geq 3$  and  $a \in \{2,3\}$ . as follows: contains at least two  $K_n$ .

$$q(p) = 1.$$

$$q(v_i^j) = \begin{cases} 2; i = 1, & j = 3. \\ i + 1; 1 \leq i \leq n - 1, & j = 1. \\ i + 2; 1 \leq i \leq n - 1, & j = 2; \\ 2 \leq i \leq n - 1, & j = 3. \end{cases}$$

Then the code of vertices are:

$$q_{\Pi}(p) = \begin{cases} 0, & 1^{st} \text{ tuple.} \\ 1, & \text{tuple other.} \end{cases}$$

$$q_{\Pi}(v_i^j) = \begin{cases} 0, & 2^{nd} \text{ tuple; } i = 1 \text{ and } j = 3; \\ i + 1^{th} \text{ tuple; } 1 \leq i \leq n - 1 \text{ and } j = 1; \\ i + 2^{th} \text{ tuple; } 1 \leq i \leq n - 1 \text{ and } j = 2; \\ i + 2^{th} \text{ tuple; } 2 \leq i \leq n - 1 \text{ and } j = 3. \\ 2, & 2^{nd} \text{ tuple; } 1 \leq i \leq n - 1 \text{ and } j = 2; \\ 3^{rd} \text{ tuple; } 1 \leq i \leq n - 1 \text{ and } j = 3; \\ n + 1^{th} \text{ tuple; } 1 \leq i \leq n - 1 \text{ and } j = 1. \\ 1, & \text{otherwise.} \end{cases}$$

Since all vertices in  $aK_n$  for  $n \geq 3$  and  $a \in \{2,3\}$  have distinct color codes, then  $q$  is a locating coloring. So,  $\chi_L(aK_n) \leq n + 1$ . Therefore  $\chi_L(aK_n) = n + 1$ . ■

**Theorem 3.2.** *The locating chromatic number for amalgamation of complete graph  $(4K_n)$ ,*

$$\chi_L(4K_n) = \begin{cases} 5; & \text{if } n = 3, \\ n + 1; & \text{if } n \geq 4. \end{cases}$$

**Proof.**

Consider two cases.

**Case 1 ( $n = 3$ ).**

The graph  $4K_3$  contains  $K_3$ , based on Theorem 1.1, we have  $\chi_L(4K_3) \geq 3$ . Suppose that there is a 3-locating coloring of  $4K_3$ . Without loss of generality, we assign  $q(p) = 1$ , color 2 or 3 to other vertices. It is clearly, there are two vertices have the same color codes. So,  $\chi_L(4K_3) \geq 4$ . Suppose that there is a 4 locating of  $4K_3$ . Without loss of generality, we assign  $q(p) = 1$ , color 2, 3 or 4 to other vertices. Then there are two pairs vertices with the same color such that have the same color codes, a contrary. As a result,  $\chi_L(4K_3) \geq 5$ .

Next, vertex coloring is given to determine the locating chromatic number of the graph. Let  $q$  be a vertex coloring that uses 5 colors. The  $4K_3$  graph is given a color class such that it is obtained

$$Q_1 = \{p\}, Q_2 = \{v_1^1, v_1^3\}, Q_3 = \{v_2^1, v_1^2, v_1^4\}, Q_4 = \{v_2^2, v_2^3\}, Q_5 = \{v_2^4\}.$$

Then, the color codes of vertices are:

$$\begin{aligned} q_\pi(p) &= \{0,1,1,1,1\}; & q_\pi(v_1^1) &= \{1,0,1,2,2\}; & q_\pi(v_2^1) &= \{1,1,0,2,2\}; \\ q_\pi(v_1^2) &= \{1,2,0,1,2\}; & q_\pi(v_2^2) &= \{1,2,1,0,2\}; & q_\pi(v_1^3) &= \{1,0,2,1,2\}; \\ q_\pi(v_2^3) &= \{1,1,2,0,2\}; & q_\pi(v_1^4) &= \{1,2,0,2,1\}; & q_\pi(v_2^4) &= \{1,2,1,2,0\}. \end{aligned}$$

Thus, since all vertices in  $4K_3$  have distinct color codes, then  $q$  is a 5-locating coloring. So,  $\chi_L(4K_3) \geq 5$ . Therefore, we have  $\chi_L(4K_3) = 5$ .

**Case 2 ( $n \geq 4$ ).**

We will first determine the lower bound of the locating chromatic number of the graph  $4K_n$ , Since  $4K_n$  contains at least two  $K_n$  and based on Theorem 1.1, then  $\chi_L(aK_n) \geq n + 1$ . Consequently,  $\chi_L(4K_n) \geq n + 1$ .

Next, to determine the upper bound of the locating chromatic number of  $4K_n$  for  $n \geq 4$ . Let  $q$  be a coloring using  $(n + 1)$  colors as follows:

$$q(p) = 1.$$

$$q(v_i^j) = \begin{cases} 2; i = 1, j = 3. \\ i + 1; 1 \leq i \leq n - 1, & j = 1; \\ & i = 1, 2, & j = 4. \\ i + 2; 1 \leq i \leq n - 1, & j = 2; \\ & 2 \leq i \leq n - 1, & j = 3; \\ & 3 \leq i \leq n - 1, & j = 4. \end{cases}$$

Then, the color code of vertices are:

$$q_{\Pi}(p) = \begin{cases} 0, & 1^{st} \text{ tuple} \\ 1, & \text{tuple other} \end{cases}$$

$$q_{\Pi}(v_i^j) = \begin{cases} 0, & 2^{nd} \text{ tuple; } i = 1 \text{ and } j = 3; \\ & i + 1^{th} \text{ tuple; } 1 \leq i \leq n - 1 \text{ and } j = 1; \\ & i + 1^{th} \text{ tuple; } i = 1, 2 \text{ and } j = 4; \\ & i + 2^{th} \text{ tuple; } 1 \leq i \leq n - 1 \text{ and } j = 2; \\ & i + 2^{th} \text{ tuple; } 2 \leq i \leq n - 1 \text{ and } j = 3; \\ & i + 2^{th} \text{ tuple; } 3 \leq i \leq n - 1 \text{ and } j = 4. \\ 2, & 2^{nd} \text{ tuple; } 1 \leq i \leq n - 1 \text{ and } j = 2; \\ & 3^{rd} \text{ tuple; } 1 \leq i \leq n - 1 \text{ and } j = 3; \\ & 4^{th} \text{ tuple; } 1 \leq i \leq n - 1 \text{ and } j = 4; \\ & n + 1^{th} \text{ tuple; } 1 \leq i \leq n - 1 \text{ and } j = 1. \\ 1, & \text{otherwise.} \end{cases}$$

Since all vertices in  $4K_n$  for  $n \geq 4$  have distinct color codes, then  $q$  is a locating coloring. So,  $\chi_L(4K_n) \leq n + 1$ . Therefore  $\chi_L(4K_n) = n + 1$ . ■

**Theorem 3.3.** *The locating chromatic number of amalgamation of complete graph  $(5K_n)$ ,*

$$\chi_L(5K_n) = \begin{cases} n + 2; & \text{if } n = 3, 4, \\ n + 1; & \text{if } n \geq 5. \end{cases}$$

**Proof.**

Consider two cases.

**Case 1.** Consider the following two subcases.

**Subcase 1.1 ( $n = 3$ )**

We will first determine the lower bound of the locating chromatic number of the graph  $5K_3$  because the graph  $5K_3$  contains  $K_3$ ; based on Theorem 1.1, we obtain  $\chi_L(5K_3) \geq 3$ . Suppose that there is a 3 – locating coloring of  $4K_3$ . Without loss of generality, we assign  $q(p) = 1$ , then the vertices neighboring  $p$  are given the colors 2 or 3. It is clearly that there are two vertices have the same color code, a contrary. So,  $\chi_L(5K_3) \geq 4$ . Similarly, if we use 4 colors, as a result, we have  $\chi_L(5K_3) \geq 5$ .

Next, vertex coloring is given to determine the locating chromatic number of the graph. Let  $q$  be a vertex coloring that uses 5 colors. The  $5K_3$  graph is given a color class such that it is obtained

$$Q_1 = \{p\}, Q_2 = \{v_1^1, v_1^3\}, Q_3 = \{v_2^1, v_1^2, v_1^4\}, Q_4 = \{v_2^2, v_2^3, v_1^5\}, Q_5 = \{v_2^4, v_2^5\}.$$

Then, the color codes of vertices are:

$$\begin{aligned} q_\pi(p) &= \{0,1,1,1,1\}; & q_\pi(v_1^1) &= \{1,0,1,2,2\}; & q_\pi(v_2^1) &= \{1,1,0,2,2\}; \\ q_\pi(v_1^2) &= \{1,2,0,1,2\}; & q_\pi(v_2^2) &= \{1,2,1,0,2\}; & q_\pi(v_1^3) &= \{1,0,2,1,2\}; \\ q_\pi(v_2^3) &= \{1,1,2,0,2\}; & q_\pi(v_1^4) &= \{1,2,0,2,1\}; & q_\pi(v_2^4) &= \{1,2,1,2,0\}; \\ q_\pi(v_1^5) &= \{1,2,2,0,1\}; & q_\pi(v_2^5) &= \{1,2,2,1,0\}. \end{aligned}$$

Thus, since all vertices in  $5K_3$  have distinct color codes, then  $q$  is a 5-locating coloring. So,  $\chi_L(5K_3) \geq 5$ . Therefore, we have  $\chi_L(5K_3) = 5$ .

**Subcase 1.2 (n = 4)**

We will first determine the lower bound of the locating chromatic number of the graph  $5K_4$ . Since the graph  $5K_4$  contains  $K_4$ , based on Theorem 1.1, we obtain  $\chi_L(5K_4) \geq 4$ . Suppose that there is a 4 – locating coloring of  $5K_4$ . Without loss of generality, we assign  $q(p) = 1$ , then the vertices neighboring  $p$  are given the colors 2,3 or 4. It is clear that there are two vertices that have the same color codes, a contrary. So,  $\chi_L(5K_4) \geq 5$ . Similarly, if we use 5 colors. As a results, we have  $\chi_L(5K_4) \geq 6$ .

Let  $q$  be a vertex coloring that uses 6 colors. The  $5K_4$  graph is given a color class such that it is obtained

$$\begin{aligned} Q_1 &= \{p\}, Q_2 = \{v_1^1, v_1^3, v_1^4\}, Q_3 = \{v_2^1, v_1^2, v_2^4, v_1^5\}, Q_4 = \{v_3^1, v_2^2, v_2^3, v_2^5\}, \\ Q_5 &= \{v_3^2, v_3^3, v_3^4\}, Q_6 = \{v_3^5\}. \end{aligned}$$

Then, the color codes of vertices are:

$$\begin{aligned} q_\pi(p) &= \{0,1,1,1,1,1\}; & q_\pi(v_1^1) &= \{1,0,1,1,2,2\}; & q_\pi(v_2^1) &= \{1,1,0,1,2,2\}; \\ q_\pi(v_3^1) &= \{1,1,1,0,2,2\}; & q_\pi(v_1^2) &= \{1,2,0,1,1,2\}; & q_\pi(v_2^2) &= \{1,2,1,0,1,2\}; \\ q_\pi(v_3^2) &= \{1,2,1,1,0,2\}; & q_\pi(v_1^3) &= \{1,0,2,1,1,2\}; & q_\pi(v_2^3) &= \{1,1,2,0,1,2\}; \\ q_\pi(v_3^3) &= \{1,1,2,1,0,2\}; & q_\pi(v_1^4) &= \{1,0,1,2,1,2\}; & q_\pi(v_2^4) &= \{1,1,0,2,1,2\}; \\ q_\pi(v_3^4) &= \{1,1,1,2,0,2\}. \end{aligned}$$

Thus, since all vertices in  $5K_4$  have distinct color code, then  $q$  is a 6-locating coloring. So,  $\chi_L(5K_4) \geq 6$ . Therefore, we have  $\chi_L(5K_4) = 6$ .

**Case 2 ( $n \geq 5$ ).**

We will first determine the lower bound of the locating chromatic number of the graph  $5K_n$ . Since  $5K_n$  contains at least two  $K_n$  and based on Theorem 1.1, then  $\chi_L(5K_n) \geq n + 1$ .

Next, to determine the upper bound of the locating chromatic number of  $5K_n$  for  $n \geq 5$ . Let  $q$  be a coloring using  $(n + 1)$  colors as follows:

$$q(p) = 1.$$

$$q(v_i^j) = \begin{cases} 2; i = 1, j = 3 \\ i + 1; 1 \leq i \leq n - 1, & j = 1 \\ i = 1, 2, 3, & j = 5 \\ i + 2; 1 \leq i \leq n - 1, & j = 2 \\ 2 \leq i \leq n - 1, & j = 3 \\ 4 \leq i \leq n - 1, & j = 5 \end{cases}$$

Then, the color codes of vertices are:

$$q_{\Pi}(p) = \begin{cases} 0, 1^{st} \text{ tuple} \\ 1, \text{ tuple other} \end{cases}$$

$$q_{\Pi}(v_i^j) = \begin{cases} 0, 2^{nd} \text{ tuple; } i = 1 \text{ and } j = 3; \\ i + 1^{th} \text{ tuple; } 1 \leq i \leq n - 1 \text{ and } j = 1; \\ i + 1^{th} \text{ tuple; } i = 1, 2 \text{ and } j = 4; \\ i + 1^{th} \text{ tuple; } i = 1, 2, 3 \text{ and } j = 5; \\ i + 2^{th} \text{ tuple; } 1 \leq i \leq n - 1 \text{ and } j = 2; \\ i + 2^{th} \text{ tuple; } 2 \leq i \leq n - 1 \text{ and } j = 3; \\ i + 2^{th} \text{ tuple; } 3 \leq i \leq n - 1 \text{ and } j = 4; \\ n + 1^{th} \text{ tuple; } i = 4 \text{ and } j = 5. \\ 2, 2^{nd} \text{ tuple; } 1 \leq i \leq n - 1 \text{ and } j = 2; \\ 3^{rd} \text{ tuple; } 1 \leq i \leq n - 1 \text{ and } j = 3; \\ 4^{th} \text{ tuple; } 1 \leq i \leq n - 1 \text{ and } j = 4; \\ 5^{th} \text{ tuple; } 1 \leq i \leq n - 1 \text{ and } j = 5; \\ n + 1^{th} \text{ tuple; } 1 \leq i \leq n - 1 \text{ and } j = 1. \\ 1, \text{ otherwise.} \end{cases}$$

Since all vertices in  $5K_n$  for  $n \geq 5$  have distinct color codes, then  $q$  is a locating coloring. So,  $\chi_L(5K_n) \leq n + 1$ . Therefore  $\chi_L(5K_n) = n + 1$ . ■

**Theorem 3.4.** *The locating chromatic number of amalgamation of complete graph  $6K_n$ ,*

$$\chi_L(6K_n) = \begin{cases} n + 2; \text{ if } n = 3, 4, 5 \\ n + 1; \text{ if } n \geq 6. \end{cases}$$

**Proof.**

Consider two cases.

**Case 1.** Consider the following three subcases.

**Subcase 1.1 ( $n = 3$ )**

We will first determine the lower bound of the locating chromatic number of the graph  $6K_3$ . Since the graph  $6K_3$  contains  $K_3$ , based on Theorem 1.1, we obtain  $\chi_L(6K_3) \geq 3$ . Suppose that there is a 3 – locating coloring of  $6K_3$ . Without loss of generality, we assign  $q(p) = 1$ , then the vertices neighboring  $p$  are given the colors 2 or 3. It is clear that two vertices that have the same color codes, a contrary. So,  $\chi_L(6K_3) \geq 4$ . Similarly, if we use 5 colors, as a result, we have  $\chi_L(6K_3) \geq 5$ .

Next, vertex coloring is given to determine the locating chromatic number of the graph. Let  $q$  be a vertex coloring that uses 5 colors. The  $6K_3$  graph is given a color class such that it is obtained

$$Q_1 = \{p\}, Q_2 = \{v_1^1, v_1^3, v_1^5\}, Q_3 = \{v_2^1, v_1^2, v_1^4\}, Q_4 = \{v_2^2, v_2^3, v_2^6\}, Q_5 = \{v_2^4, v_2^5, v_2^6\}.$$

Then, the color codes of vertices are:

$$\begin{aligned} q_\pi(p) &= \{0,1,1,1,1\}; & q_\pi(v_1^1) &= \{1,0,1,2,2\}; & q_\pi(v_2^1) &= \{1,1,0,2,2\}; \\ q_\pi(v_1^2) &= \{1,2,0,1,2\}; & q_\pi(v_2^2) &= \{1,2,1,0,2\}; & q_\pi(v_1^3) &= \{1,0,2,1,2\}; \\ q_\pi(v_2^3) &= \{1,1,2,0,2\}; & q_\pi(v_1^4) &= \{1,2,0,2,1\}; & q_\pi(v_2^4) &= \{1,2,1,2,0\}; \\ q_\pi(v_1^5) &= \{1,0,2,2,1\}; & q_\pi(v_2^5) &= \{1,1,2,2,0\}; & q_\pi(v_1^6) &= \{1,2,2,0,1\}; \\ q_\pi(v_2^6) &= \{1,2,2,1,0\}. \end{aligned}$$

Thus, since all vertices in  $6K_3$  have distinct color code, then  $q$  is a 5-locating coloring. So,  $\chi_L(6K_3) \geq 5$ . Therefore, we have  $\chi_L(6K_3) = 5$ .

**Subcase 1.2 ( $n = 4$ )**

We will first determine the lower bound of the locating chromatic number of the graph  $6K_4$ . Since the graph  $6K_4$  contains  $K_4$ , based on Theorem 1.1, we obtain  $\chi_L(6K_4) \geq 4$ . Without loss of generality, we assign  $q(p) = 1$ , and then the vertices neighboring  $p$  are given the colors 2,3,3 or 4. It is clear that two vertices have the same color codes, a contrary. So,  $\chi_L(6K_4) \geq 5$ . Similarly, if we use 5 color, subsequently,  $\chi_L(6K_4) \geq 6$ .

Next, vertex coloring is given to determine the locating chromatic number of the graph. Let  $q$  be a vertex coloring that uses 6 colors. The  $6K_4$  graph is given a color class such that it is obtained

$$\begin{aligned} Q_1 &= \{p\}, Q_2 = \{v_1^1, v_1^3, v_1^4\}, Q_3 = \{v_2^1, v_1^2, v_2^4, v_1^5, v_1^6\}, Q_4 = \{v_3^1, v_2^2, v_2^3, v_2^5\}, \\ Q_5 &= \{v_3^2, v_3^3, v_3^4, v_2^6\}, Q_6 = \{v_3^5, v_3^6\}. \end{aligned}$$

Then, the color codes of vertices are:

$$\begin{aligned} q_\pi(p) &= \{0,1,1,1,1,1\}; & q_\pi(v_1^1) &= \{1,0,1,1,2,2\}; & q_\pi(v_2^1) &= \{1,1,0,1,2,2\}; \\ q_\pi(v_3^1) &= \{1,1,1,0,2,2\}; & q_\pi(v_2^2) &= \{1,2,0,1,1,2\}; & q_\pi(v_2^3) &= \{1,2,1,0,1,2\}; \end{aligned}$$



$$\begin{aligned}
 q_\pi(v_3^2) &= \{1,2,1,1,0,2\}; & q_\pi(v_1^3) &= \{1,0,2,1,1,2\}; & q_\pi(v_2^3) &= \{1,1,2,0,1,2\}; \\
 q_\pi(v_3^3) &= \{1,1,2,1,0,2\}; & q_\pi(v_1^4) &= \{1,0,1,2,1,2\}; & q_\pi(v_2^4) &= \{1,1,0,2,1,2\}; \\
 q_\pi(v_3^4) &= \{1,1,1,2,0,2\}; & q_\pi(v_1^5) &= \{1,2,0,1,2,1\}; & q_\pi(v_2^5) &= \{1,2,1,0,2,1\}; \\
 q_\pi(v_3^5) &= \{1,2,1,1,2,0\}; & q_\pi(v_1^6) &= \{1,2,0,2,1,1\}; & q_\pi(v_2^6) &= \{1,2,1,2,0,1\}; \\
 q_\pi(v_3^6) &= \{1,2,1,2,1,0\}.
 \end{aligned}$$

Thus, since all vertices in  $6K_4$  have distinct color codes,  $q$  is a 6-locating coloring. So,  $\chi_L(6K_4) \geq 6$ . Therefore, we have  $\chi_L(6K_4) = 6$ .

**Subcase 1.3 (n = 5)**

We will first determine the lower bound of the locating chromatic number of the graph  $6K_5$ . Since the graph  $6K_5$  contains  $K_5$ , based on Theorem 1.1, we obtain  $\chi_L(6K_5) \geq 5$ . Without loss of generality, we assign  $q(p) = 1$ , and then the vertices neighboring  $p$  are given the colors 2,3,4 or 5. It is clear that two vertices have the same color codes, a contrary. So,  $\chi_L(6K_5) \geq 6$ . Similarly, if we use 6 colors, thus,  $\chi_L(6K_5) \geq 7$ .

Next, vertex coloring is given to determine the locating chromatic number of the graph. Let  $q$  be a vertex coloring that uses 7 colors. The  $6K_5$  graph is given a color class such that it is obtained

$$\begin{aligned}
 Q_1 &= \{p\}, Q_2 = \{v_1^1, v_1^3, v_1^4, v_1^5\}, Q_3 = \{v_2^1, v_1^2, v_2^4, v_2^5, v_1^6\}, Q_4 = \{v_3^1, v_2^2, v_2^3, v_3^5, v_2^6\}, \\
 Q_5 &= \{v_4^1, v_2^2, v_3^3, v_3^4, v_3^6\}, Q_6 = \{v_4^2, v_4^3, v_4^4, v_4^5\}, Q_7 = \{v_4^6\}.
 \end{aligned}$$

Then, the color codes of vertices are:

$$\begin{aligned}
 q_\pi(p) &= \{0,1,1,1,1,1\}; & q_\pi(v_1^1) &= \{1,0,1,1,1,2,2\}; & q_\pi(v_2^1) &= \{1,1,0,1,1,2,2\}; \\
 q_\pi(v_3^1) &= \{1,1,1,0,1,2,2\}; & q_\pi(v_4^1) &= \{1,1,1,1,0,2,2\}; & q_\pi(v_1^2) &= \{1,2,0,1,1,1,2\}; \\
 q_\pi(v_2^2) &= \{1,2,1,0,1,1,2\}; & q_\pi(v_3^2) &= \{1,2,1,1,0,1,2\}; & q_\pi(v_4^2) &= \{1,2,1,1,1,0,2\}; \\
 q_\pi(v_1^3) &= \{1,0,2,1,1,1,2\}; & q_\pi(v_2^3) &= \{1,1,2,0,1,1,2\}; & q_\pi(v_3^3) &= \{1,1,2,1,0,1,2\}; \\
 q_\pi(v_4^3) &= \{1,1,2,1,1,0,2\}; & q_\pi(v_1^4) &= \{1,0,1,2,1,1,2\}; & q_\pi(v_2^4) &= \{1,1,0,2,1,1,2\}; \\
 q_\pi(v_3^4) &= \{1,1,1,2,0,1,2\}; & q_\pi(v_4^4) &= \{1,1,1,2,1,0,2\}; & q_\pi(v_1^5) &= \{1,0,1,1,2,1,2\}; \\
 q_\pi(v_2^5) &= \{1,1,0,1,2,1,2\}; & q_\pi(v_3^5) &= \{1,1,1,0,2,1,2\}; & q_\pi(v_4^5) &= \{1,1,1,1,2,0,2\}; \\
 q_\pi(v_1^6) &= \{1,2,0,1,1,2,1\}; & q_\pi(v_2^6) &= \{1,2,1,0,1,2,1\}; & q_\pi(v_3^6) &= \{1,2,1,1,0,2,1\}; \\
 q_\pi(v_4^6) &= \{1,2,1,1,1,2,0\}.
 \end{aligned}$$

Thus, since all vertices in  $6K_5$  have distinct color codes,  $q$  is a 7-locating coloring. So,  $\chi_L(6K_5) \geq 7$ . Therefore, we have  $\chi_L(6K_5) = 7$ .

**Case 2 ( $n \geq 6$ ).**

We will first determine the lower bound of the locating chromatic number of the graph  $6K_n$ . Since  $6K_n$  contains at least two  $K_n$  based on Theorem 1.1, then  $\chi_L(6K_n) \geq n + 1$ .

Next, to determine the upper bound of the locating chromatic number of  $6K_n$  for  $n \geq 6$ . Let  $q$  be a coloring using  $(n + 1)$  colors as follows:

$$q(p) = 1.$$

$$q(v_i^j) = \begin{cases} 2; i = 1, j = 3 \\ 7; i = 5, j = 6 \\ i + 1; 1 \leq i \leq n - 1, j = 1 \\ i = 1, 2, j = 4 \\ i = 1, 2, 3, j = 5 \\ i = 1, 2, 3, 4, j = 6 \\ i + 2; 1 \leq i \leq n - 1, j = 2 \\ 2 \leq i \leq n - 1, j = 3 \\ 3 \leq i \leq n - 1, j = 4 \\ 4 \leq i \leq n - 1, j = 5 \end{cases}$$

Then, the color code of vertices are:

$$q_{\Pi}(p) = \begin{cases} 0, 1^{st} \text{ tuple.} \\ 1, \text{ otherwise.} \end{cases}$$

$$q_{\Pi}(v_i^j) = \begin{cases} 0, 2^{nd} \text{ tuple; } i = 1 \text{ and } j = 3; \\ i + 1^{th} \text{ tuple; } 1 \leq i \leq n - 1 \text{ and } j = 1; \\ i + 1^{th} \text{ tuple; } 1 \leq i \leq n - 2 \text{ and } j = 6; \\ i + 1^{th} \text{ tuple; } 1 \leq i \leq n - 3 \text{ and } j = 5; \\ i + 1^{th} \text{ tuple; } 1 \leq i \leq n - 4 \text{ and } j = 4; \\ i + 2^{th} \text{ tuple; } 1 \leq i \leq n - 1 \text{ and } j = 2; \\ i + 2^{th} \text{ tuple; } 2 \leq i \leq n - 1 \text{ and } j = 3; \\ i + 2^{th} \text{ tuple; } 3 \leq i \leq n - 1 \text{ and } j = 4; \\ i + 2^{th} \text{ tuple; } 4 \leq i \leq n - 1 \text{ and } j = 5; \\ n + 2^{th} \text{ tuple; } i = 5 \text{ and } j = 6. \\ 2, n + 1^{th} \text{ tuple; } 1 \leq i \leq n - 1 \text{ and } j = 1; \\ 2^{nd} \text{ tuple; } 1 \leq i \leq n - 1 \text{ and } j = 2; \\ 3^{rd} \text{ tuple; } 1 \leq i \leq n - 1 \text{ and } j = 3; \\ 4^{th} \text{ tuple; } 1 \leq i \leq n - 1 \text{ and } j = 4; \\ 5^{th} \text{ tuple; } 1 \leq i \leq n - 1 \text{ and } j = 5; \\ 6^{th} \text{ tuple; } 1 \leq i \leq n - 1 \text{ and } j = 6. \\ 1, \text{ otherwise.} \end{cases}$$

Since all vertices in  $6K_n$  for  $n \geq 6$  have distinct color codes,  $q$  is a locating coloring. So,  $\chi_L(6K_n) \leq n + 1$ . Therefore  $\chi_L(6K_n) = n + 1$ . ■

Figure 2 is an example of the locating chromatic number for the amalgamation of complete graph  $4K_4$ , with the locating chromatic number being 5.




Figure 2. A minimum locating coloring of  $4K_4$

## ACKNOWLEDGEMENTS

The research was supported by the Directorate of Higher Education, Ministry of Education and Culture. The authors are thankful for the funding of the research

## REFERENCES

- [1] P. J. Slater, "Leaves of Trees," in *Proceedings of the Sixth Southeastern Conference on Combinatorics, Graph Theory and Computing*, Boca Raton, Florida: Utilitas Mathematica Publishing, Inc., 1975, pp. 549–559.
- [2] G. Chartrand, E. Salehi, and P. Zhang, "The Partition Dimension of a Graph," *Aequationes Math*, vol. 59, 2000.
- [3] G. Chartrand, D. Erwin, M. A. Henning, P. J. Slater, and P. Zhang, "Locating Chromatic Number of a Graph," *Bull. Inst. Combin. Appl.*, vol. 36, pp. 89–101, 2002.
- [4] G. Chartrand, D. Erwin, M. A. Henning, P. J. Slater, and P. Zhang, "Graphs of order  $n$  with locating-chromatic number  $n - 1$ ," *Discrete Math*, vol. 269, no. 1–3, pp. 65–79, Jul. 2003, doi: 10.1016/S0012-365X(02)00829-4.
- [5] H. Assiyatun and E. Baskoro, "Locating-Chromatic Number of Amalgamation of Stars," 2011.
- [6] E. T. Baskoro, H. Assiyatun, D. Suprijanto, R. Simanjuntak, and S. Uttunggadewa, "The Locating-Chromatic Number of Firecracker Graphs," *Far East Journal of Mathematical Sciences (FJMS)*, vol. 63, no. 1, pp. 11–23, 2012, [Online]. Available: <http://pphmj.com/journals/fjms.htm>
- [7] Asmiati and E. T. Baskoro, "Characterizing all graphs containing cycles with locating-chromatic number 3," in *AIP Conference Proceedings*, 2012, pp. 351–357. doi: 10.1063/1.4724167.

- [8] E. T. Baskoro and Asmiati, “Characterizing all trees with locating-chromatic number 3,” *Electronic Journal of Graph Theory and Applications*, vol. 1, no. 2, pp. 109–117, 2013, [Online]. Available: [www.ejgta.org](http://www.ejgta.org)
- [9] Asmiati, “On the locating-chromatic numbers of non-homogeneous caterpillars and firecracker graphs,” *Far East Journal of Mathematical Sciences*, vol. 100, no. 8, pp. 1305–1316, Oct. 2016, doi: 10.17654/MS100081305.
- [10] Asmiati, I. K. Sadha Gunce Yana, and L. Yulianti, “On the Locating Chromatic Number of Certain Barbell Graphs,” *Int J Math Math Sci*, vol. 2018, 2018, doi: 10.1155/2018/5327504.
- [11] Asmiati, A. Irawan, A. Nuryaman, and K. Muludi, “The locating chromatic number for certain operation of origami graphs,” *Mathematics and Statistics*, vol. 11, no. 1, pp. 101–106, Jan. 2023, doi: 10.13189/ms.2023.110111.
- [12] A. Asmiati, M. Damayanti, and L. Yulianti, “On the Locating Chromatic Number of Barbell Shadow Path Graph,” *Indonesian Journal of Combinatorics*, vol. 5, no. 2, p. 82, Dec. 2021, doi: 10.19184/ijc.2021.5.2.4.
- [13] Asmiati, L. Yulianti, Aldino, Aristoteles, and A. Junaidi, “The Locating Chromatic Number of a Disjoint Union of Some Double Stars,” in *Journal of Physics: Conference Series*, Institute of Physics Publishing, Oct. 2019. doi: 10.1088/1742-6596/1338/1/012035.
- [14] A. Irawan and Asmiati, “The locating-chromatic number of subdivision firecracker graphs,” *International Mathematical Forum*, vol. 13, no. 10, pp. 485–492, 2018, doi: 10.12988/imf.2018.8844.
- [15] A. Irawan, Asmiati, Suharsono, and K. Muludi, “The Locating-Chromatic Number for Certain Operation of Generalized Petersen Graphs  $SP(4,2)$ ,” in *Journal of Physics: Conference Series*, Institute of Physics Publishing, Oct. 2019. doi: 10.1088/1742-6596/1338/1/012033.
- [16] A. Irawan, Asmiati, S. Suharsono, K. Muludi, and L. Zakaria, “Certain Operation of Generalized Petersen Graphs having Locating-Chromatic Number Five,” *Advances and Applications in Discrete Mathematics*, vol. 24, no. 2, pp. 83–97, Jul. 2020, doi: 10.17654/dm024020083.
- [17] A. Irawan, A. Asmiati, L. Zakaria, and K. Muludi, “The Locating-Chromatic Number of Origami Graphs,” *Algorithms*, vol. 14, no. 6, 2021, doi: 10.3390/a14060167.
- [18] A. Irawan, B. Herindri Samodra Utami, A. Nuryaman, and K. Muludi, “A Procedure for Determining The Locating Chromatic Number of an Origami Graphs,” *IJCSNS International Journal of Computer Science and Network Security*, vol. 22, no. 9, 2022, doi: 10.22937/IJCSNS.2022.22.9.5.
- [19] Asmiati, W. Okzarima, Notiragayu, and L. Zakaria, “Upper Bounds of the Locating Chromatic Numbers of Shadow Cycle Graphs,” *Int J Math Comput Sci*, vol. 19, no. 1, pp. 239–248, 2024, [Online]. Available: <http://ijmcs.future-in-tech.net>
- [20] A. Behtoei and B. Omoomi, “On the locating chromatic number of Kneser graphs,” *Discrete Appl Math*, vol. 159, no. 18, pp. 2214–2221, Dec. 2011, doi: 10.1016/j.dam.2011.07.015.
- [21] A. Behtoei and B. Omoomi, “On the Locating Chromatic Number of the Cartesian Product of Graphs,” Jun. 2011, [Online]. Available: <http://arxiv.org/abs/1106.3453>
- [22] M. Ghanem, H. Al-Ezeh, and A. Dabbour, “Locating Chromatic Number of Powers of Paths and Cycles,” *Symmetry (Basel)*, vol. 11, Mar. 2019, doi: 10.3390/sym11030389.
- [23] M. Furuya and N. Matsumoto, “Upper bounds on the locating chromatic number of trees,” *Discrete Appl Math*, vol. 257, pp. 338–341, Mar. 2019, doi: 10.1016/j.dam.2018.10.034.

- [24] K. Prawinasti, M. Ansori, Asmiati, Notiragayu, and A. R. G N Rofi, “The Locating Chromatic Number for Split Graph of Cycle,” in *Journal of Physics: Conference Series*, IOP Publishing Ltd, Jan. 2021. doi: 10.1088/1742-6596/1751/1/012009.
- [25] M. Damayanti, Asmiati, Fitriani, M. Ansori, and A. Faradilla, “The Locating Chromatic Number of some Modified Path with Cycle having Locating Number Four,” in *Journal of Physics: Conference Series*, IOP Publishing Ltd, Jan. 2021. doi: 10.1088/1742-6596/1751/1/012008.
- [26] S. Rahmatalia, A. Asmiati, and N. Notiragayu, “Bilangan Kromatik Lokasi Graf Split Lintasan,” *Jurnal Matematika Integratif*, vol. 18, no. 1, p. 73, May 2022, doi: 10.24198/jmi.v18.n1.36091.73-80.