

# On Randers Change of a Generalized Exponential Metric

Meera Mishra\* and R. K. Pandey

Department of Mathematics & Computer Science, BBD University-U.P., India

Email: \*meera.mishra@bbdu.ac.in

## Abstract

In this paper, we study the properties of a special  $(\alpha, \beta)$ -metric  $e^{k_1 \frac{\beta}{\alpha}} + \beta e^{k_2 \frac{\beta}{\alpha}}$ , the Randers change of the generalized exponential metric. We find the necessary and sufficient condition for this metric to be locally projectively flat and we also prove the conditions for this metric to be of the Berwald and Douglas type.

**Keywords:** Berwald space; Douglas space; Finsler space;  $(\alpha, \beta)$ -metric; Projectively flat.

## Abstrak

*Pada artikel ini akan dipelajari sifat-sifat khusus dari  $(\alpha, \beta)$ -metric  $e^{k_1 \frac{\beta}{\alpha}} + \beta e^{k_2 \frac{\beta}{\alpha}}$ , perubahan Randers dari metrik eksponensial umum. Kami menemukan syarat perlu dan cukup agar metrik ini menjadi datar secara lokal dan kami juga membuktikan syarat agar metrik ini bertipe Berwald dan Douglas.*

**Kata Kunci:** Ruang Berwald; Ruang Douglas; Ruang Finsler;  $(\alpha, \beta)$ -metric; Projectively flat.

2020MSC: 53B20.

## 1. INTRODUCTION

Graph In 1972, M. Matsumoto [1] [2] introduced the notion of the  $(\alpha, \beta)$ -metric on the basis of the Randers metric. There are several types of  $(\alpha, \beta)$ -metrics, such as the Randers metric  $\alpha + \beta$  [3], Kropina metric  $\frac{\alpha^2}{\beta}$  [1] [4], generalized Kropina metric  $\frac{\alpha^{m+1}}{\beta^m}$  ( $m \neq 0, -1$ ) [5] and Matsumoto metric  $\frac{\alpha^2}{\alpha - \beta}$  [4]. In these metrics,  $\alpha(x, y) = (a_{ij}(x)y^i y^j)^{1/2}$  is a Riemannian metric and  $\beta(x, y) = b_i(x)y^i$  is a differential one form [6].

In 2013, G. Shanker and Ravindra [7] introduced a special  $(\alpha, \beta)$ -metric  $\alpha e^{\beta/\alpha} + \beta$  and considered it a Randers change in the exponential metric. Now, we generalize this concept by introducing an  $(\alpha, \beta)$ -metric

$$F(\alpha, \beta) = \alpha e^{k_1 \frac{\beta}{\alpha}} + \beta e^{k_2 \frac{\beta}{\alpha}}, \quad (1)$$

where  $k_1$  and  $k_2$  are some constants and are referred to as the Randers changes of the generalized exponential metric.

In this paper, we have studied the properties of this  $(\alpha, \beta)$ -metric, identified the necessary and sufficient conditions for the metric to be locally projectively flat and determined the conditions for this metric to be of the Berwald and Douglas type. A deep study of the projectively flat Finsler metric was also carried out by authors [5] [8] [9] [10] [11] [12] and [13].

\* Corresponding author

Submitted August 13<sup>th</sup>, 2024, Revised October 25<sup>th</sup>, 2024,

Accepted for publication November 15<sup>th</sup>, 2024, Published Online November 30<sup>th</sup>, 2024

©2024 The Author(s). This is an open-access article under CC-BY-SA license (<https://creativecommons.org/licence/by-sa/4.0/>)

## 2. PRELIMINARIES

Let  $\alpha = (a_{ij}(x)y^i y^j)^{1/2}$  be a Riemannian metric,  $\beta = b_i y^i$  be a 1-form and  $F = \alpha\phi(s)$ , where  $s = \frac{\beta}{\alpha}$ ,  $\phi = \phi(s)$  be a positive  $C^\infty$  function defined in a neighborhood of the origin  $s = 0$ . Since we know that  $F = \alpha\phi\left(\frac{\beta}{\alpha}\right)$  is a Finsler metric for any  $\alpha$  and  $\beta$  with  $b = \|\beta\|_\alpha < b_0$  iff

$$\phi(s) > 0, \phi(s) - s\phi'(s) + (b^2 - s^2)\phi''(s) > 0, (|s| \leq b < b_0).$$

Replacing  $b$  by  $s$ , we obtain

$$\phi(s) - s\phi'(s) > 0, (|s| < b_0).$$

Let  $G^i$  and  $G_\alpha^i$  denote the spray coefficients of  $F$  and  $\alpha$  respectively, defined as

$$G^i = \frac{g^{il}}{4} \{ [F^2]_{x^k y^l} y^k - [F^2]_{x^k} \}, G_\alpha^i = \frac{a^{il}}{4} \{ [\alpha^2]_{x^k y^l} y^k - [\alpha^2]_{x^k} \},$$

where  $g_{ij} = \frac{1}{2} [F^2]_{y^i y^j}$  and

$$(a^{ij}) = (a_{ij})^{-1}, F_{x^k} = \frac{\partial F}{\partial x^k}, F_{y^k} = \frac{\partial F}{\partial y^k}.$$

For an  $(\alpha, \beta)$ -metric  $L(\alpha, \beta)$ , the space  $\mathbb{R}^n = (M^n, \alpha)$  is known as the Riemannian space associated with the Finsler space  $F^n = (M^n, L(\alpha, \beta))$ . The covariant differentiation with respect to the Levi-Civita connection  $\gamma_{jk}^i(x)$  of  $\mathbb{R}^n$  is denoted by  $(;)$ . Now, we have the following [14]:

**Lemma 2.1.** The spray coefficients  $G^i$  are related to  $G_\alpha^i$  by

$$G^i = G_\alpha^i + \alpha Q s_0^i + J(-2\alpha Q s_0 + r_{00}) \frac{y^i}{\alpha} + H(-2\alpha Q s_0 + r_{00}) \left\{ b^i - \frac{y^i}{\alpha} \right\}, \quad (2)$$

where

$$\begin{aligned} Q &= \frac{\phi'}{\phi - s\phi'}, \\ J &= \frac{(\phi - s\phi')\phi'}{2\phi((\phi - s\phi') + (b^2 - s^2)\phi'')}, \\ H &= \frac{\phi''}{2((\phi - s\phi') + (b^2 - s^2)\phi'')} \end{aligned}$$

where  $s_{l0} = s_{li} y^i$ ,  $s_0 = s_{l0} b^l$ ,  $r_{00} = r_{ij} y^i y^j$ ,  $r_{ij} = \frac{1}{2}(b_{i;j} + b_{j;i})$ ,  $s_{ij} = \frac{1}{2}(b_{i;j} - b_{j;i})$ ,  $r_j^i = a^{ir} r_{rj}$ ,  $s_j^i = a^{ir} s_{rj}$ ,  $r_j = b_r r_j^r$ ,  $s_j = b_r s_j^r$ ,  $b^i = a^{ir} b_r$  and  $b^2 = a^{rs} b_r b_s$ . It is well-known that [15] a Finsler metric  $F = F(x, y)$  on an open subset  $U \subset \mathbb{R}^n$  is projectively flat if and only if

$$F_{x^k y^l} y^k - F_{x^l} = 0. \quad (3)$$

By equation (3), we have the following lemma [9].

**Lemma 2.2.** An  $(\alpha, \beta)$ -metric  $F = \alpha\phi(s)$ , where  $s = \frac{\beta}{\alpha}$ , is projectively flat on an open subset  $U \subset \mathbb{R}^n$  if and only if

$$(a_{ml}\alpha^2 - y_m y_l)G_\alpha^m + \alpha^3 Qs_{l0} + H\alpha(-2\alpha Qs_0 + r_{00})(b_l\alpha - sy_l) = 0. \quad (4)$$

The functions  $G^i(x, y)$  of  $F^n$  with an  $(\alpha, \beta)$ -metric can be written in the form [16]

$$2G^i = \gamma_{00}^i + 2B^i, \quad (5)$$

$$B^i = \frac{\alpha L_\beta}{L_\alpha} s_0^i + C^* \left\{ \frac{\beta L_\beta}{\alpha L} y^i - \frac{\alpha L_{\alpha\alpha}}{L_\alpha} \left( \frac{1}{\alpha} y^i - \frac{\alpha}{\beta} b^i \right) \right\}. \quad (6)$$

provided  $\beta^2 + L_\alpha + \alpha\gamma^2 L_{\alpha\alpha} \neq 0$ , where  $\gamma^2 = b^2\alpha^2 - \beta^2$ ,  $L_\alpha = \frac{\partial L}{\partial \alpha}$ ,  $L_\beta = \frac{\partial L}{\partial \beta}$ ,  $L_{\alpha\alpha} = \frac{\partial^2 L}{\partial \alpha^2}$ , and  $s_0^i = s_j^i y^j$ . Now we put

$$C^* = \frac{\alpha\beta(r_{00}L_\alpha - 2\alpha s_0 L_\beta)}{2(\beta^2 L_\alpha + \alpha\gamma^2 L_{\alpha\alpha})}.$$

We denote the homogeneous polynomials in  $(y^i)$  of degree  $r$  by  $hp(r)$  for brevity. For example,  $\gamma_{00}^i$  is  $hp(2)$ . From equation (5), the Berwald connection  $B\Gamma = (G_{jk}^i, G_j^i, 0)$  of  $F^n$  with an  $(\alpha, \beta)$ -metric is given by [16]

$$\begin{aligned} G_j^i &= \partial_j G^i = \gamma_{0j}^i + B_j^i \\ G_{jk}^i &= \partial_k G_j^i = \gamma_{jk}^i + B_{jk}^i, \end{aligned}$$

where  $B_j^i = \partial_j B^i$  and  $B_{jk}^i = \partial_k B_j^i$ . According to [16],  $B_{jk}^i$  are determined by

$$L_\alpha B_{ji}^k y^j y_t + \alpha L_\beta (B_{ji}^k b_t - b_{j,i}) y^j = 0, \quad (7)$$

where  $y_k = a_{ik} y^i$ . A Finsler space  $F^n$  with an  $(\alpha, \beta)$ -metric is a Douglas space if and only if  $B^{ij} \equiv B^i y^j - b^j y^i$  is  $hp(3)$  [17]. From equation (6),  $B^{ij}$  can be written as follows:

$$B^{ij} = \frac{\alpha L_\beta}{L_\alpha} (s_0^i y^j - s_0^j y^i) + \frac{\alpha^2 L_{\alpha\alpha}}{\beta L_\alpha} C^* (b^i y^j - b^j y^i). \quad (8)$$

### 3. RESULTS AND DISCUSSION

#### 3.1. Projectively flat $(\alpha, \beta)$ -metric

In this context, we consider the metric  $F = \alpha e^{k_1 \frac{\beta}{\alpha}} + \beta e^{k_2 \frac{\beta}{\alpha}}$  which is obtained by the Randers change of the generalized exponential metric

$$F = \alpha\phi(s), \phi(s) = (e^{k_1 s} + s e^{k_2 s}), s = \frac{\beta}{\alpha}. \quad (9)$$

Let  $b_0 > 0$  be the largest number satisfying

$$\phi(s) - s\phi'(s) + (b^2 - s^2)\phi''(s) > 0, (|s| \leq b < b_0),$$

that is,

$$e^{k_1 s}(1 - sk_1 + b^2 k_1^2 - s^2 k_1^2) + e^{k_2 s}(b^2 sk_2^2 - s^2 k_2 + 2bk_2 - s^3 k_2^2 - 2k_2 s^2) > 0, (|s| \leq b < b_0).$$

**Lemma 3.1.**  $F = \alpha e^{k_1 \frac{\beta}{\alpha}} + \beta e^{k_2 \frac{\beta}{\alpha}}$  is a Finsler metric, iff  $\|\beta\|_\alpha < 1$ .

**Proof.**

If  $F = \alpha e^{k_1 \frac{\beta}{\alpha}} + \beta e^{k_2 \frac{\beta}{\alpha}}$  is a Finsler metric, then

$$e^{k_1 s}(1 - sk_1 + b^2 k_1^2 - s^2 k_1^2) + e^{k_2 s}(b^2 sk_2^2 - s^2 k_2 + 2bk_2 - s^3 k_2^2 - 2k_2 s^2) > 0, (|s| \leq b < b_0).$$

Let  $s = b$ , then we obtain  $b < 1, \forall b < b_0$ . Let  $b \rightarrow b_0$ , then  $b_0 < 1$ . Therefore  $\|\beta\|_\alpha < 1$ . Now, if  $|s| \leq b < 1$  then

$$e^{k_1 s}(1 - sk_1 + b^2 k_1^2 - s^2 k_1^2) + e^{k_2 s}(b^2 sk_2^2 - s^2 k_2 + 2bk_2 - s^3 k_2^2 - 2k_2 s^2) \geq e^{k_1 s}(1 - sk_1) + e^{k_2 s}(2bk_2 - 3k_2 s^2) > 0 \quad (\text{since } b^2 - s^2 \geq 0).$$

Thus  $F = \alpha e^{k_1 \frac{\beta}{\alpha}} + \beta e^{k_2 \frac{\beta}{\alpha}}$  is a Finsler metric. By Lemma (2.1), the spray coefficients  $G^i$  of  $F$  are written by equation (2) as

$$\begin{aligned} Q &= \frac{k_1 e^{k_1 s} + k_2 s e^{k_2 s} + e^{k_2 s}}{e^{k_1 s} - k_1 s e^{k_1 s} - k_2 s^2 e^{k_2 s}}, \quad s = \frac{\beta}{\alpha}, \\ J &= \frac{(1 - sk_1)k_1 e^{2k_1 s} + \{(1 - sk_1)sk_2 + (1 - sk_1) - s^2 k_1 k_2\}e^{(k_1 + k_2)s} - s^2 k_2(1 + sk_2)e^{2k_2 s}}{2(e^{k_1 s} + s e^{k_2 s})\{(1 - sk_1) + (b^2 - s^2)k_1\}e^{k_1 s} + [(b^2 - s^2)k_2^2 s + 2b^2 k_2 - 3s^2 k_2]e^{k_2 s}}, \\ H &= \frac{k_1^2 e^{k_1 s} + k_2^2 s e^{k_2 s} + 2k_2 e^{k_2 s}}{2\{(1 - sk_1) + (b^2 - s^2)k_1\}e^{k_1 s} + [(b^2 - s^2)k_2^2 s + 2b^2 k_2 - 3s^2 k_2]e^{k_2 s}}. \end{aligned}$$

Then equation (4) is reduced to the following form:

$$\begin{aligned} (a_{ml}\alpha^2 - y_m y_l)G_\alpha^m + \alpha^3 \frac{k_1 e^{k_1 s} + k_2 s e^{k_2 s} + e^{k_2 s}}{e^{k_1 s} - k_1 s e^{k_1 s} - k_2 s^2 e^{k_2 s}} s_{l0} + \\ \frac{\alpha(k_1^2 e^{k_1 s} + k_2^2 s e^{k_2 s} + 2k_2 e^{k_2 s})}{2\{(1 - sk_1) + (b^2 - s^2)k_1\}e^{k_1 s} + [(b^2 - s^2)k_2^2 s + 2b^2 k_2 - 3s^2 k_2]e^{k_2 s}} \left[ -2\alpha \frac{(k_1 e^{k_1 s} + k_2 s e^{k_2 s} + e^{k_2 s})}{(e^{k_1 s} - k_1 s e^{k_1 s} - k_2 s^2 e^{k_2 s})} s_0 + r_{00} \right] \times \\ (b_l \alpha - s y_l) = 0, s = \frac{\beta}{\alpha}. \end{aligned} \quad (10)$$

Now we use the following result [18]:

**Lemma 3.2.** If  $(a_{ml}\alpha^2 - y_m y_l)G_\alpha^m = 0$ , then  $\alpha$  is projectively flat.

**Proof.**

If  $(a_{ml}\alpha^2 - y_m y_l)G_\alpha^m = 0$ , then we have

$$\alpha^2 a_{ml} G_\alpha^m = y_m G_\alpha^m y_l.$$

Hence, if there is an  $\eta = \eta(x, y)$  such that  $y_m G_\alpha^m = \alpha^2 \eta$ , we find

$$a_{ml} G_\alpha^m = \eta y_l. \quad (11)$$

Contracting equation (11) with  $a^{il}$  yields  $G_\alpha^i = \eta y^i$ , and hence  $\alpha$  is projectively flat.

**Theorem 3.3.** The  $(\alpha, \beta)$ -metric  $F = \alpha e^{k_1 \frac{\beta}{\alpha}} + \beta e^{k_2 \frac{\beta}{\alpha}}$  is locally projectively flat iff  $\beta$  is parallel with respect to  $\alpha$  and  $\alpha$  is locally projectively flat, i.e., of constant curvature.

**Proof.**

Suppose, that  $F$  is locally projectively flat. First, we rewrite equation (10) as a polynomial in  $y^i$  and  $\alpha$ , then we obtain

$$\begin{aligned} & \{-2\alpha^2 \beta [(1 + k_1) + b^2] + 2\beta^3 k_1\} e^{k_1 \frac{\beta}{\alpha}} (a_{ml} \alpha^2 - y_m y_l) G_\alpha^m + (2\alpha^6 + 2b^2 \alpha^6 - 2\alpha^4 \beta^2) \left[ k_1 e^{k_1 \frac{\beta}{\alpha}} + \right. \\ & \left. (1 + k_2 \frac{\beta}{\alpha}) e^{k_2 \frac{\beta}{\alpha}} \right] s_{l0} - (2\alpha^4 s_0 \left[ k_1 e^{k_1 \frac{\beta}{\alpha}} + \left(1 + k_2 \frac{\beta}{\alpha}\right) e^{k_2 \frac{\beta}{\alpha}} \right] - \alpha^2 \beta e^{k_1 \frac{\beta}{\alpha}} r_{00}) (b_l \alpha^2 - \beta y_l) \} + \alpha \{ (2\alpha^2 + 2b^2 \alpha^2) e^{k_1 \frac{\beta}{\alpha}} (a_{ml} \alpha^2 - y_m y_l) G_\alpha^m - 2\alpha^4 \beta \left[ k_1 e^{k_1 \frac{\beta}{\alpha}} + \left(1 + k_2 \frac{\beta}{\alpha}\right) e^{k_2 \frac{\beta}{\alpha}} \right] s_{l0} + \right. \\ & \left. \alpha^2 e^{k_1 \frac{\beta}{\alpha}} r_{00} (b_l \alpha^2 - \beta y_l) - k_2 \beta^2 r_{00} e^{k_2 \frac{\beta}{\alpha}} (b_l \alpha^2 - \beta y_l) \} = 0, \\ & \text{or } U + \alpha V = 0, \end{aligned} \quad (12)$$

where

$$\begin{aligned} U = & \{-2\alpha^2 \beta [(1 + k_1) + b^2] + 2\beta^3 k_1\} e^{k_1 \frac{\beta}{\alpha}} (a_{ml} \alpha^2 - y_m y_l) G_\alpha^m + (2\alpha^6 + 2b^2 \alpha^6 - 2\alpha^4 \beta^2) \times \\ & \left[ k_1 e^{k_1 \frac{\beta}{\alpha}} + \left(1 + k_2 \frac{\beta}{\alpha}\right) e^{k_2 \frac{\beta}{\alpha}} \right] s_{l0} - (2\alpha^4 s_0 \left[ k_1 e^{k_1 \frac{\beta}{\alpha}} + \left(1 + k_2 \frac{\beta}{\alpha}\right) e^{k_2 \frac{\beta}{\alpha}} \right] + \alpha^2 \beta e^{k_1 \frac{\beta}{\alpha}} r_{00}) (b_l \alpha^2 - \beta y_l), \end{aligned}$$

and

$$\begin{aligned} V = & 2\alpha^2 (1 + b^2) e^{k_1 \frac{\beta}{\alpha}} (a_{ml} \alpha^2 - y_m y_l) G_\alpha^m - 2\alpha^4 \beta \left[ k_1 e^{k_1 \frac{\beta}{\alpha}} + \left(1 + k_2 \frac{\beta}{\alpha}\right) e^{k_2 \frac{\beta}{\alpha}} \right] s_{l0} + \\ & (\alpha^2 e^{k_1 \frac{\beta}{\alpha}} - \beta^2 k_2 e^{k_2 \frac{\beta}{\alpha}}) r_{00} (b_l \alpha^2 - \beta y_l). \end{aligned}$$

Now, equation (12) is a polynomial in  $y^i$ , such that  $U$  and  $V$  are rational in  $y^i$  and  $\alpha$  is irrational. Therefore, we must have  $U = 0$  and  $V = 0$  which implies that

$$\begin{aligned} & \{-2\alpha^2 \beta [(1 + k_1) + b^2] + 2\beta^3 k_1\} e^{k_1 \frac{\beta}{\alpha}} (a_{ml} \alpha^2 - y_m y_l) G_\alpha^m = \left( 2\alpha^4 s_0 \left[ k_1 e^{k_1 \frac{\beta}{\alpha}} + \left(1 + k_2 \frac{\beta}{\alpha}\right) e^{k_2 \frac{\beta}{\alpha}} \right] + \alpha^2 \beta e^{k_1 \frac{\beta}{\alpha}} r_{00} \right) \times \\ & (b_l \alpha^2 - \beta y_l) - (2\alpha^6 + 2b^2 \alpha^6 - 2\alpha^4 \beta^2) \left[ k_1 e^{k_1 \frac{\beta}{\alpha}} + \left(1 + k_2 \frac{\beta}{\alpha}\right) e^{k_2 \frac{\beta}{\alpha}} \right] s_{l0}, \end{aligned} \quad (13)$$

and

$$\begin{aligned} & 2\alpha^2 (1 + b^2) e^{k_1 \frac{\beta}{\alpha}} (a_{ml} \alpha^2 - y_m y_l) G_\alpha^m - 2\alpha^4 \beta \left[ k_1 e^{k_1 \frac{\beta}{\alpha}} + \left(1 + k_2 \frac{\beta}{\alpha}\right) e^{k_2 \frac{\beta}{\alpha}} \right] s_{l0} - \\ & (\alpha^2 e^{k_1 \frac{\beta}{\alpha}} - \beta^2 k_2 e^{k_2 \frac{\beta}{\alpha}}) r_{00} (b_l \alpha^2 - \beta y_l), \end{aligned} \quad (14)$$

respectively. Contracting equations (13) and (3.6) with  $b^l$ , we have

$$\begin{aligned} & \{-2\alpha^2\beta[1+k_1+b^2]+2\beta^3k_1\}e^{k_1\frac{\beta}{\alpha}}(b_m\alpha^2-y_m\beta)G_\alpha^m \\ & \quad = -2\alpha^6s_0\left[k_1e^{k_1\frac{\beta}{\alpha}}+\left(1+k_2\frac{\beta}{\alpha}\right)e^{k_2\frac{\beta}{\alpha}}\right]+\alpha^2\beta e^{k_1\frac{\beta}{\alpha}}r_{00}(b^2\alpha^2-\beta^2) \\ & 2\alpha^2(1+b^2)e^{k_1\frac{\beta}{\alpha}}(b_m\alpha^2-y_m\beta)G_\alpha^m \\ & \quad = 2\alpha^4\beta\left[k_1e^{k_1\frac{\beta}{\alpha}}+\left(1+k_2\frac{\beta}{\alpha}\right)e^{k_2\frac{\beta}{\alpha}}\right]s_0-(\alpha^2e^{k_1\frac{\beta}{\alpha}}-\beta^2k_2e^{k_2\frac{\beta}{\alpha}})r_{00}(b^2\alpha^2-\beta^2) \end{aligned} \quad (15)$$

$$\begin{aligned} & 2(1+b^2)e^{k_1\frac{\beta}{\alpha}}(b_m\alpha^2-y_m\beta)G_\alpha^m \\ & \quad = 2\alpha^2\beta\left[k_1e^{k_1\frac{\beta}{\alpha}}+\left(1+k_2\frac{\beta}{\alpha}\right)e^{k_2\frac{\beta}{\alpha}}\right]s_0-\left(e^{k_1\frac{\beta}{\alpha}}-\frac{\beta^2}{\alpha^2}k_2e^{k_2\frac{\beta}{\alpha}}\right)r_{00}(b^2\alpha^2-\beta^2) \end{aligned} \quad (16)$$

Multiplying equation (16) by  $\alpha^2\beta$  and adding the result obtained to equation (15), we get

$$\begin{aligned} & \beta\left[2k_1(-\alpha^2+\beta^2)(b_m\alpha^2-y_m\beta)e^{k_1\frac{\beta}{\alpha}}G_\alpha^m-k_2\beta^2r_{00}e^{k_2\frac{\beta}{\alpha}}(b^2\alpha^2-\beta^2)\right] \\ & \quad = 2\alpha^4(-\alpha^2+\beta^2)\left[k_1e^{k_1\frac{\beta}{\alpha}}+\left(1+k_2\frac{\beta}{\alpha}\right)e^{k_2\frac{\beta}{\alpha}}\right]s_0 \end{aligned} \quad (17)$$

The polynomial  $2\alpha^4(-\alpha^2+\beta^2)\left[k_1e^{k_1\frac{\beta}{\alpha}}+\left(1+k_2\frac{\beta}{\alpha}\right)e^{k_2\frac{\beta}{\alpha}}\right]$  is not divisible by  $\beta$  and  $\beta$  is not divisible by

$$2\alpha^4(-\alpha^2+\beta^2)\left[k_1e^{k_1\frac{\beta}{\alpha}}+\left(1+k_2\frac{\beta}{\alpha}\right)e^{k_2\frac{\beta}{\alpha}}\right].$$

Thus  $s_0$  is divisible by  $\beta$  and  $[2k_1(-\alpha^2+\beta^2)(b_m\alpha^2-y_m\beta)e^{k_1\frac{\beta}{\alpha}}G_\alpha^m-k_2\beta^2r_{00}e^{k_2\frac{\beta}{\alpha}}(b^2\alpha^2-\beta^2)]$  is divisible by  $2\alpha^4(-\alpha^2+\beta^2)\left[k_1e^{k_1\frac{\beta}{\alpha}}+\left(1+k_2\frac{\beta}{\alpha}\right)e^{k_2\frac{\beta}{\alpha}}\right]$ . Hence, there exist scalar functions  $\tau = \tau(x), \chi = \chi(x)$  such that

$$s_0 = \tau\beta, \quad (18)$$

$$\begin{aligned} & 2k_1(-\alpha^2+\beta^2)(b_m\alpha^2-y_m\beta)e^{k_1\frac{\beta}{\alpha}}G_\alpha^m-k_2\beta^2r_{00}e^{k_2\frac{\beta}{\alpha}}(b^2\alpha^2-\beta^2) \\ & \quad = \chi\left\{2\alpha^4(-\alpha^2+\beta^2)\left[k_1e^{k_1\frac{\beta}{\alpha}}+\left(1+k_2\frac{\beta}{\alpha}\right)e^{k_2\frac{\beta}{\alpha}}\right]\right\}. \end{aligned} \quad (19)$$

Then, equation (17) reduces to

$$\beta\chi\left\{2\alpha^4(-\alpha^2+\beta^2)\left[k_1e^{k_1\frac{\beta}{\alpha}}+\left(1+k_2\frac{\beta}{\alpha}\right)e^{k_2\frac{\beta}{\alpha}}\right]\right\}=2\alpha^4(-\alpha^2+\beta^2)\left[k_1e^{k_1\frac{\beta}{\alpha}}+\left(1+k_2\frac{\beta}{\alpha}\right)e^{k_2\frac{\beta}{\alpha}}\right]\tau\beta.$$

Thus  $\tau = \chi$ . Hence, equation (15) becomes

$$\begin{aligned} & \chi\left\{2\alpha^4(-\alpha^2+\beta^2)\left[k_1e^{k_1\frac{\beta}{\alpha}}+\left(1+k_2\frac{\beta}{\alpha}\right)e^{k_2\frac{\beta}{\alpha}}\right]\right\}\{-2\alpha^2\beta[1+k_1+b^2]+2\beta^3k_1+2k_1\beta\alpha^2\}= \\ & \quad \left\{(2\alpha^2\beta[1+k_1+b^2]-2\beta^3k_1)k_2\beta^2e^{k_2\frac{\beta}{\alpha}}+\alpha^2\beta e^{k_1\frac{\beta}{\alpha}}(-\alpha^2+\beta^2)2k_1\right\}r_{00}(b^2\alpha^2-\beta^2). \end{aligned} \quad (20)$$

Since  $(b^2\alpha^2 - \beta^2)$  is not divisible by

$$\frac{\left\{2\alpha^4(-\alpha^2+\beta^2)\left[k_1e^{k_1\frac{\beta}{\alpha}}+(1+k_2\frac{\beta}{\alpha})e^{k_2\frac{\beta}{\alpha}}\right]\right\}\{-2\alpha^2\beta[1+k_1+b^2]+2\beta^3k_1+2k_1\beta\alpha^2\}}{\left\{(2\alpha^2\beta[1+k_1+b^2]-2\beta^3k_1)k_2\beta^2e^{k_2\frac{\beta}{\alpha}}+\alpha^2\beta e^{k_1\frac{\beta}{\alpha}}(-\alpha^2+\beta^2)2k_1\right\}},$$

it follows from (20) that  $\chi = 0$ . By (18), (19) and (20), we get

$$s_0 = 0, \quad (21)$$

$$r_{00} = 0, \quad (22)$$

$$\text{and } 2k_1(-\alpha^2 + \beta^2)(b_m\alpha^2 - y_m\beta)e^{k_1\frac{\beta}{\alpha}}G_\alpha^m - k_2\beta^2r_{00}e^{k_2\frac{\beta}{\alpha}}(b^2\alpha^2 - \beta^2) = 0.$$

Using equation (22) in the above equation, we obtain

$$(b_m\alpha^2 - y_m\beta)e^{k_1\frac{\beta}{\alpha}}G_\alpha^m = 0. \quad (23)$$

Substituting equations(22) and (23) into (14), we find

$$s_{l0} = 0. \quad (24)$$

Using equation (23) and lemma (3.2), gives  $\alpha$  is projectively flat. Using equations (22) and (24), we get  $b_{i,j} = 0$ . i.e,  $\beta$  is parallel with respect to  $\alpha$ . Conversely, if  $\beta$  is parallel with respect to  $\alpha$  and  $\alpha$  is locally projectively flat, then by Lemma (2.2), we can easily see that  $F$  is locally projectively flat.

### 3.2. Berwald and Douglas Spaces

In this part, we find the condition that a Finsler space  $F^n$  with the  $(\alpha, \beta)$ -metric (9) is a Berwald space. In  $n$ -dimensional Finsler space  $F^n$  with the  $(\alpha, \beta)$ -metric (9), we have

$$\begin{aligned} L_\alpha &= \frac{(\alpha - k_1\beta)e^{k_1\frac{\beta}{\alpha}}}{\alpha}, \quad L_\beta = k_1e^{k_1\frac{\beta}{\alpha}} + e^{k_2\frac{\beta}{\alpha}}\left(1 + k_2\frac{\beta}{\alpha}\right), \\ L_{\alpha\alpha} &= \frac{1}{\alpha}\left(\frac{k_1\beta}{\alpha}\right)^2 e^{k_1\frac{\beta}{\alpha}}, \quad L_{\beta\beta} = \frac{k_1^2}{\alpha}e^{k_1\frac{\beta}{\alpha}} + \frac{k_2}{\alpha}\left(\frac{2\alpha + k_2\beta}{\alpha}\right)e^{k_2\frac{\beta}{\alpha}} \end{aligned} \quad (25)$$

Substituting (25) in (7), we find

$$\alpha e^{k_1\frac{\beta}{\alpha}}B_{ji}^ty^jy_t - k_1\beta e^{k_1\frac{\beta}{\alpha}}B_{ji}^ty^jy_t + \left\{\alpha^2\left(k_1e^{k_1\frac{\beta}{\alpha}} + e^{k_2\frac{\beta}{\alpha}}\right) + \beta\alpha k_2e^{k_2\frac{\beta}{\alpha}}\right\}(B_{ji}^tb_t - b_{j,i})y^j = 0. \quad (26)$$

Let  $F^n$  be a Berwald space, i.e.,  $G_{jk}^i = G_{jk}^i(x)$ . Then we obtain  $B_{ji}^t = B_{ji}^t(x)$ . Since  $\alpha$  is irrational in  $(y^i)$ , hence from equation (26), we can write

$$e^{k_1\frac{\beta}{\alpha}}B_{ji}^ty^jy_t = 0, \quad (27)$$

$$-k_1\beta e^{k_1\frac{\beta}{\alpha}}B_{ji}^ty^jy_t + \left\{\alpha^2\left(k_1e^{k_1\frac{\beta}{\alpha}} + e^{k_2\frac{\beta}{\alpha}}\right) + \beta\alpha k_2e^{k_2\frac{\beta}{\alpha}}\right\}(B_{ji}^tb_t - b_{j,i})y^j = 0. \quad (28)$$

Using equations (27) and (28), we obtain

$$B_{ji}^t y^j y_t = 0 \text{ and } \left\{ \alpha^2 \left( k_1 e^{k_1 \frac{\beta}{\alpha}} + e^{k_2 \frac{\beta}{\alpha}} \right) + \beta \alpha k_2 e^{k_2 \frac{\beta}{\alpha}} \right\} (B_{ji}^t b_t - b_{j,i}) y^j = 0,$$

which gives  $B_{ji}^t a_{th} + B_{hi}^t a_{tj} = 0, B_{ji}^t b_t - b_{j,i} = 0$ . Thus, by the well known Christoffel process we find  $B_{ji}^t = 0$ . Therefore we have:

**Theorem 4.1.** The Randers change of the generalized exponential metric (9) provides a Berwald metric if and only if  $b_{j,i} = 0$ , and then the Berwald connection is Riemannian  $(\gamma_{jk}^i, \gamma_{0j}^i, 0)$ .

Now, we consider the condition for a Finsler space  $F^n$  with the  $(\alpha, \beta)$ -metric (9) to be a Douglas space. Substituting equation (25) into equation (2.7), we get

$$\begin{aligned} & 2(\alpha - k_1 \beta) e^{k_1 \frac{\beta}{\alpha}} [\alpha^2 - k_1 \alpha \beta + (b^2 \alpha^2 - \beta^2) k_1^2] B^{ij} \\ & - 2\alpha \left[ \alpha k_1 e^{k_1 \frac{\beta}{\alpha}} + (\alpha + \beta k_2) e^{k_2 \frac{\beta}{\alpha}} \right] [\alpha^2 - k_1 \alpha \beta + (b^2 \alpha^2 - \beta^2) k_1^2] (s_0^i y^j - s_0^j y^i) \\ & - k_1 \alpha^2 \left\{ (\alpha - k_1 \beta) e^{k_1 \frac{\beta}{\alpha}} r_{00} - 2\alpha s_0 \left[ \alpha k_1 e^{k_1 \frac{\beta}{\alpha}} + (\alpha + \beta k_2) e^{k_2 \frac{\beta}{\alpha}} \right] \right\} (b^i y^j - b^j y^i) = 0. \end{aligned} \quad (29)$$

Let  $F^n$  be a Douglas space, i.e.,  $B^{ij}$  are  $hp(3)$ . Separating equation (29) in rational and irrational terms of  $y^i$ , because  $\alpha$  is irrational in  $(y^i)$ , we find

$$\begin{aligned} & \left[ (-4k_1 \alpha^2 \beta + 2k_1^3 \beta^3 - 2k_1^3 b^2 \alpha^2 \beta) e^{k_1 \frac{\beta}{\alpha}} B^{ij} \right. \\ & + \left( \alpha k_1 e^{k_1 \frac{\beta}{\alpha}} + \alpha e^{k_2 \frac{\beta}{\alpha}} + \beta k_2 e^{k_2 \frac{\beta}{\alpha}} \right) (-2\alpha^3 - 2k_1^2 b^2 \alpha^3 + 2k_1^2 \alpha \beta^2) (s_0^i y^j - s_0^j y^i) \\ & + k_1^2 \alpha^2 \beta e^{k_1 \frac{\beta}{\alpha}} r_{00} (b^i y^j - b^j y^i) + 2k_1 \alpha^3 s_0 \left( \alpha k_1 e^{k_1 \frac{\beta}{\alpha}} + \alpha e^{k_2 \frac{\beta}{\alpha}} + \beta k_2 e^{k_2 \frac{\beta}{\alpha}} \right) (b^i y^j - b^j y^i) \left. \right] \\ & + \alpha \left[ e^{k_1 \frac{\beta}{\alpha}} B^{ij} (2\alpha^2 + 2k_1^2 b^2 \alpha^2) + 2k_1 \alpha \beta \left( \alpha k_1 e^{k_1 \frac{\beta}{\alpha}} + \alpha e^{k_2 \frac{\beta}{\alpha}} + \beta k_2 e^{k_2 \frac{\beta}{\alpha}} \right) (s_0^i y^j - s_0^j y^i) \right. \\ & \left. - k_1 \alpha^2 e^{k_1 \frac{\beta}{\alpha}} r_{00} (b^i y^j - b^j y^i) \right] = 0. \end{aligned} \quad (30)$$

Hence equation (30) is divided into the following two equations:

$$\begin{aligned} & \left[ (-4k_1 \alpha^2 \beta + 2k_1^3 \beta^3 - 2k_1^3 b^2 \alpha^2 \beta) e^{k_1 \frac{\beta}{\alpha}} B^{ij} \right. \\ & + \left( \alpha k_1 e^{k_1 \frac{\beta}{\alpha}} + \alpha e^{k_2 \frac{\beta}{\alpha}} + \beta k_2 e^{k_2 \frac{\beta}{\alpha}} \right) (-2\alpha^3 - 2k_1^2 b^2 \alpha^3 + 2k_1^2 \alpha \beta^2) (s_0^i y^j - s_0^j y^i) \\ & + k_1^2 \alpha^2 \beta e^{k_1 \frac{\beta}{\alpha}} r_{00} (b^i y^j - b^j y^i) + 2k_1 \alpha^3 s_0 \left( \alpha k_1 e^{k_1 \frac{\beta}{\alpha}} + \alpha e^{k_2 \frac{\beta}{\alpha}} + \beta k_2 e^{k_2 \frac{\beta}{\alpha}} \right) (b^i y^j - b^j y^i) \left. \right] = 0, \end{aligned} \quad (31)$$

and



$$e^{k_1 \frac{\beta}{\alpha}} B^{ij} (2\alpha^2 + 2k_1^2 b^2 \alpha^2) + 2k_1 \alpha \beta \left( \alpha k_1 e^{k_1 \frac{\beta}{\alpha}} + \alpha e^{k_2 \frac{\beta}{\alpha}} + \beta k_2 e^{k_2 \frac{\beta}{\alpha}} \right) (s_0^i y^j - s_0^j y^i) - k_1 \alpha^2 e^{k_1 \frac{\beta}{\alpha}} r_{00} (b^i y^j - b^j y^i) = 0. \quad (32)$$

Eliminating  $B^{ij}$  from equations (31) and (32), we obtain

$$A(s_0^i y^j - s_0^j y^i) + B(b^i y^j - b^j y^i) = 0, \quad (33)$$

where

$$A = 2\alpha(-2\alpha^4 - 4k_1^2 b^2 \alpha^4 + 6k_1^2 \alpha^2 \beta^2 - 2k_1^4 b^4 \alpha^4 + 4k_1^4 b^2 \alpha^2 \beta^2 - 2k_1^4 \beta^4) \times \left( \alpha k_1 e^{k_1 \frac{\beta}{\alpha}} + \alpha e^{k_2 \frac{\beta}{\alpha}} + \beta k_2 e^{k_2 \frac{\beta}{\alpha}} \right), \quad (34)$$

$$B = 2\alpha[k_1^2 \alpha e^{k_1 \frac{\beta}{\alpha}} r_{00} (\beta^3 k_1^2 - \alpha^2 \beta) + 2k_1 \alpha^4 (1 + k_1^2 b^2) \left( \alpha k_1 e^{k_1 \frac{\beta}{\alpha}} + \alpha e^{k_2 \frac{\beta}{\alpha}} + \beta k_2 e^{k_2 \frac{\beta}{\alpha}} \right) s_0]. \quad (35)$$

Transvecting (33) by  $b_i y_j$ , we have

$$A s_0 \alpha^2 + B(b^2 \alpha^2 - \beta^2) = 0. \quad (36)$$

The term of the equation (36) which does not contain  $\alpha^2$  is  $-\beta^5 r_{00}$ . Hence there exists  $hp(5): v_5$  such that

$$-k_1^4 \beta^5 r_{00} = \alpha^2 V_5. \quad (37)$$

Now, we consider the following two cases:

- (i)  $V_5 = 0$ ,
- (ii)  $V_5 \neq 0, \alpha^2 \not\equiv 0 \pmod{\beta}$ .

Case (i). When  $V_5 = 0$ , this leads to  $r_{00} = 0$ . Therefore, substituting  $r_{00} = 0$  in equation (36), we find

$$s_0(A + \gamma^2 B_1) = 0, \quad (38)$$

where

$$B_1 = 2k_1 \alpha^2 (1 + k_1^2 b^2) \left( \alpha k_1 e^{k_1 \frac{\beta}{\alpha}} + \alpha e^{k_2 \frac{\beta}{\alpha}} + \beta k_2 e^{k_2 \frac{\beta}{\alpha}} \right).$$

If  $A + \gamma^2 B_1 = 0$ , then the term  $A + \gamma^2 B_1 = 0$  which does not contain  $\alpha^2$  is  $-4\beta^4$ . Thus there exists  $hp(2): V_2$  such that

$$-k_1^4 \beta^4 = \alpha^2 V_2.$$

Hence we have  $V_2 = 0$ , which leads to a contradiction. Therefore,  $A + \gamma^2 B_1 \neq 0$ . Hence, we get  $s_0 = 0$  from (38). Substituting  $s_0 = 0$  and  $r_{00} = 0$  into (33), we obtain

$$A(s_0^i y^j - s_0^j y^i) = 0. \quad (39)$$

If  $A = 0$ , then from equation (34), we obtain

$$2\alpha(-2\alpha^4 - 4k_1^2 b^2 \alpha^4 + 6k_1^2 \alpha^2 \beta^2 - 2k_1^4 b^4 \alpha^4 + 4k_1^4 b^2 \alpha^2 \beta^2 - 2k_1^4 \beta^4) \times$$

$$\left(\alpha k_1 e^{k_1 \frac{\beta}{\alpha}} + \alpha e^{k_2 \frac{\beta}{\alpha}} + \beta k_2 e^{k_2 \frac{\beta}{\alpha}}\right) = 0,$$

or

$$\begin{aligned} &(-2\alpha^4 - 4k_1^2 b^2 \alpha^4 + 6k_1^2 \alpha^2 \beta^2 - 2k_1^4 b^4 \alpha^4 + 4k_1^4 b^2 \alpha^2 \beta^2 - 2k_1^4 \beta^4) \times \\ &\quad \left(\alpha k_1 e^{k_1 \frac{\beta}{\alpha}} + \alpha e^{k_2 \frac{\beta}{\alpha}} + \beta k_2 e^{k_2 \frac{\beta}{\alpha}}\right) = 0. \end{aligned} \quad (40)$$

The term of the equation (40) which does not contain  $\alpha^2$  is  $-4\beta^4$ . Thus, there exists  $hp(2): V_2$  such that

$$-2k_1^4 \beta^4 = \alpha^2 V_2,$$

from which we get  $V_2 = 0$ . This is a contradiction, therefore,  $A \neq 0$ . Hence, from equation (39) we get  $s_0^i y^j - s_0^j y^i = 0$ . Now, transvecting the above equation by  $y_j$  we get  $s_0^i = 0$ , which implies  $s_{ij} = 0$ . Consequently, we find  $r_{ij} = s_{ij} = 0$ , i.e.,  $b_{i,j} = 0$ .

Case (ii). The equation (37) shows that there exists a function  $k = k(x)$  such that  $r_{00} = k(x)\alpha^2$ .

Thus, equation (36) which does not contain  $\alpha^2$  is included in the term  $-\beta^5 r_{00}$ . Hence we get  $r_{00} = 0$ . From equation (39), we get  $A(s_0^i y^j - s_0^j y^i) = 0$ . If  $A = 0$ , then it is a contradiction. Hence  $A \neq 0$ . Therefore, we obtain  $s_0^i y^j - s_0^j y^i = 0$ . Again, transvecting this equation by  $y_j$  we get  $s_0^i = 0$ . Hence, in both the cases (i) and (ii) we have  $r_{ij} = 0$  and  $s_{ij} = 0$ , i.e.,  $b_{i,j} = 0$ . Conversely if  $b_{i,j} = 0$ , then  $F^n$  is a Berwald space, so  $F^n$  is a Douglas space. Thus we have the following:

**Theorem 4.2.** The Randers change of the generalized exponential metric (9) is of the Douglas type if and only if  $\alpha^2 \not\equiv 0 \pmod{\beta}$  and  $b_{i,j} = 0$ .

From Theorem 4.1 and Theorem 4.2, we have

**Theorem 4.3.** If the Randers change in the generalized exponential metric (9) is of the Douglas type, then it is Berwaldian.

## 4. CONCLUSION

In this way we obtain the results of Randers change of the generalized Exponential metric in terms of constants  $k_1$  and  $k_2$ . By putting the different values for  $k_1$  and  $k_2$  we can find different results for their respective values.

## 5. REFERENCES

- [1] M. Hashiguchi, "On Conformal transformations of Kropina metric," *J. Math. Kyoto Univ.*, vol. 16, pp. 25-50, 1976.
- [2] M. S. Knebelman, "Conformal Geometry of generalized metric space," *Proc. Nat. Acad. Sci. USA*, vol. 15, pp. 376-379, 1929.

- [3] G. Randers, "On an asymmetrical metric in the four-space of general relativity," *Phys. Rev.*, vol. 59, p. 195, 1941.
- [4] R. S. Ingarden, "Differential Geometry and Physics," *Tensor N.S.*, vol. 30, pp. 201-209, 1976.
- [5] M. Matsumoto, "Projective changes of Finsler metric and projective flat Finsler space," *Tensor N. S.*, vol. 34, pp. 303-315, 1980.
- [6] M. Matsumoto, "A slope of a mountain is a Finsler surface with respect to time measure," *J. Math. Kyoto Univ.*, vol. 29, pp. 17-25, 1989.
- [7] G. Shanker and R. Ravindra, "On Randers change of exponential metric," *Appl. Sci.*, vol. 15, pp. 94-103, 2013.
- [8] M. Matsumoto, "Projective flat Finsler spaces with  $(\alpha, \beta)$ -metric," *Rep. Math. Phys.*, vol. 30, pp. 15-20, 1991.
- [9] H. S. Park and I. Y. Lee, "On projectively flat Finsler spaces with  $(\alpha, \beta)$ -metric," *Commun. Korean Math. Soc.*, vol. 14, no. 2, pp. 373-383, 1999.
- [10] Y. Shen and L. Zhao, "Some projectively flat  $(\alpha, \beta)$ -metrics," *Sci. China A*, vol. 49, pp. 838-851, 2006.
- [11] Z. Shen, "Projectively flat Randers metrics with constant flag curvature," *Math. Ann.*, vol. 325, pp. 19-30, 2003.
- [12] Z. Shen and G. C. Yildirim, "On a class of projectively flat metrics with constant flag curvature," *Canad. J. Math.*, vol. 60, no. 2, pp. 143-156, 2008.
- [13] Z. Shen, "On projectively flat  $(\alpha, \beta)$ -metrics," *Canad. Math. Bull.*, vol. 52, no. 1, pp. 132-144, 2009.
- [14] S. S. Chern and Z. Shen, *Riemann-Finsler Geometry*, Singapore: World Scientific, 2005.
- [15] G. Hamel, "Über die Geometrien in denen die Geraden die Kürzesten sind," *Math. Ann.*, vol. 16, pp. 25-50, 1903.
- [16] M. Matsumoto, "The Berwald connection of Finsler space with an  $(\alpha, \beta)$ -metric," *Tensor N. S.*, vol. 50, pp. 18-21, 1991.
- [17] S. Bácsó and M. Matsumoto, "On the Finsler spaces of Douglas type. A generalization of the notion of Berwald space," *Publ. Math. Debrecen*, vol. 51, pp. 385-406, 1997.
- [18] Y. Yu, "Projectively flat exponential Finsler metrics," *J. Zhejiang Univ. Sci. A*, vol. 6, pp. 1068-1076, 2006.
- [19] M. Hashiguchi, S. Hojo and M. Matsumoto, "On Landsberg spaces of two dimension with  $(\alpha, \beta)$ ," *J. Korean Math. Soc.*, vol. 10, pp. 17-26, 1973.