

A Cytotoxic Flavanone from The Pod Peels of *Tephrosia vogelii* Hook.f.

Valentina Adimurti Kusumaningtyas^{1,*}, Nia Sri Hardianti¹, Melina Melina²,
Lia Dewi Juliawaty³, Yana Maolana Syah³

¹Department of Chemistry, Faculty of Science and Informatics, Universitas Jenderal Achmad Yani,
Jalan Terusan Jenderal Sudirman, Cimahi 40285, Indonesia

²Department of Informatics, Faculty of Science and Informatics, Universitas Jenderal Achmad Yani,
Jalan Terusan Jenderal Sudirman, Cimahi 40285, Indonesia

³Natural Product Chemistry Group, Faculty of Mathematics and Natural Sciences, Institut Teknologi Bandung,
Jalan Ganesha no. 10, Bandung 40132, Indonesia

*Corresponding author: valentina.adimurti@lecture.unjani.ac.id

Received: September 2020; Revision: October 2020; Accepted: November 2020; Available online: December 2020

Abstract

Tephrosia vogelii Hook.f. is a species of the family Fabaceae (Leguminosae). These plants are termed "Polong-polongan" in Indonesia, and are known to contain active flavonoid groups. Previous studies have shown the isolation of one known flavanone: isolonchocarpin from methanol extract, and the structure obtained was established based on chemical evidence as well as spectroscopic methods, including NMR, and also by a comparison with published data. This research is aimed at evaluating the cytotoxic property of methanol extract against larvae of *Artemia salina* Leach, using the Brine Shrimp Lethality Test (BSLT) method. The results show potent cytotoxicity at LC₅₀ of 41.40 ppm.

Keyword: *Artemia salina* Leach., cytotoxic activity, Fabaceae, flavanone, *Tephrosia vogelii* Hook.f.

DOI: <https://doi.org/10.15408/jkv.v6i2.17551>

1. INTRODUCTION


Tephrosia is one of the 400 species in family Fabaceae, which belongs to the sub-family Papilionoideae, comprising shrubs, herbs and trees (Polhill, *et al.*, 1981), distributed across the tropics and sub-tropics (Tarus, *et al.*, 2002). Previous studies on the intrinsic chemical constituents showed the occurrence of chalcones (Andrei, *et al.*, 2000; Chang, *et al.*, 2000; Gomez-Garibay, *et al.*, 2002; Tarus, *et al.*, 2002), flavanones (Chang, *et al.*, 2000; Jang, *et al.*, 2003; Kishore, *et al.*, 2003), flavones (Prabhakar, *et al.*, 1996; Achmad, *et al.*, 2007), isoflavones (Yenesew, *et al.*, 1989; Dagne, *et al.*, 1992; Chang, *et al.*, 2000), rotenoid (Prashant, *et al.*, 1993; Andrei, *et al.*, 1997; Jang, *et al.*, 2003). Table 1 and Figure 1 demonstrate several highly diverse phytochemical compounds present, as

generally observed in most other similar genera, including Cassia from Fabaceae (Kusumaningtyas, *et al.*, 2020).

The phytochemical and pharmacological evaluations showed cytotoxic potentials in several compounds (Roy, *et al.*, 1986; Ganapaty, *et al.*, 2009). Touqeer, *et al.* (2013) performed similar research and reported various activities from several isolates, including cytotoxicity, as observed in Table 2 (Padmapriya, 2017). This current investigation evaluates the phytochemicals and bioactivities of compounds obtained from Indonesian *Tephrosia*. The results showed the cytotoxic effects of pod peel methanol extract from *T. vogelii* against the *Artemia salina* Leach larvae, with a Brine Shrimp Lethality Test (BSLT) method with LC₅₀ of 41.40 ppm.

Table 1. Tephrosia chemical constituents

Chemical constituents	Plant	References
Chalcones: tefon (1)	<i>T. candida</i> (seed)	Tanaka, <i>et al.</i> , 1992
Flavanones: maksimaflavanon A (2)	<i>T. maxima</i> (root)	Rao, <i>et al.</i> , 1994
Flavones: fulvinervin B (3)	<i>T. fulvinervis</i> (seed)	Rao, <i>et al.</i> , 1985
Isoflavones: kalopogoniumisoflavan B (4)	<i>T. maxima</i> (root)	Murthy, <i>et al.</i> , 1985
Rotenoid: 9-demetilidihidrostemonal (5)	<i>T. pentaphylla</i> (root)	Dagne, <i>et al.</i> , 1989

**Figure 1.** Structures of chemical constituents in *Tephrosia***Table 2.** Cytotoxic activity of *Tephrosia purpurea* extracts

Extracts	Cytotoxicity (IC ₅₀ value µg/mL)
Leaves	95.73 ± 9.60 ^{*#§}
Root	382.33 ± 18.78 [§]
Stem	324.80 ± 21.20
Seed	303.97 ± 24.31

Mean ± SD, n = 3. P < 0.05 significantly different when compared to root^{*}, stem[#] and seed[§]. (Padmapriya, *et al.*, 2017)

2. MATERIALS AND METHODS

Experimental Procedure

The melting points were obtained using a micro Fisher-John, while the NMR data were recorded with JEOL ECA 500 (¹H 500 MHz; ¹³C 125 MHz), where tetramethylsilane served as an internal standard. In addition, mass spectra were obtained on LCT XE ESI-TOF waters, and Centrifugal Thin-Layer Chromatography (Chromatotron) was performed on the silica gel 200 mesh using a precoated silica gel 60 GF₂₅₄ (0.25 mm thickness) with various mobile phases. Subsequently, the spots were visualised by 1.5% cerium sulfate in 2 N sulfuric acid spraying agent, followed by heating. The NMR spectrum was then measured using residual peaks and deuterated solvents CDCl₃.

Plant Material

The *T. vogelii* pod peels were obtained at the Plant Taxonomy Laboratory, Biology Department, FMIPA, Unpad, West Java Province, Indonesia in March 2016. These samples were identified by Drs. Joko Kusmoro, M.P. at the Jatiningor Herbarium, Indonesia, and were deposited at the herbarium with number 183/HB/03/2016.

Brine Shrimp Lethality Assay

The bioassay indicator used to monitor plant extract toxicity during fractionation involved larva *A. salina*. Moreover,

propyleneglycol/Tween 80/water (4:1:4) in 5 mL of saltwater was applied as a negative control, while ten milligrams of potassium dichromate dissolved in propyleneglycol/Tween 80/water (4:1:4) served as the positive control. The assessment procedure was as follows: ten shrimps were transferred to each sample vial containing extracts at varied concentrations of 500, 250, 125, 62.5, 31.2, 15.6, 7.8, and 0 ppm (control). Therefore artificial seawater for hatched Brine shrimp (*A. salina*) eggs, prepared from commercial sea salt 38 g/L (Sasidharan, *et al.*, 2008) was added to each sample vial to make 5 mL. This was followed by test tubes examination, and the number of dead larvae in each bottle was counted after 24 hours. The respective death percentage were determined and the test was performed triplicate. Consequently, statistical analysis used for determining the death percentage (Equation 1) and lethal concentration (LC₅₀). (Sahgal, *et al.*, 2010).

$$\%PD = \frac{(Tn - An)}{(Tn)} \times 100\% \dots\dots\dots(1)$$

Where PD is percentage of death, Tn is total nauplii and An is alive nauplii. In addition, probit analysis was performed after obtaining mortality rate. This evaluation was conducted to calculate the LC₅₀, defined as the concentration required for a compound to produced 50% death. Subsequently, the results were counted by using linear regression equation $y = a + bx$, and the statistical analysis (Melina, *et al.*, 2020) were expressed as the mean value \pm standard error of mean (SEM), while the variance was evaluated through ANOVA test. The extract toxicity level was organized according to Meyer (1982) classification. In addition, LC₅₀ scores in the range ≤ 30 ppm were defined as highly toxic, while ≤ 1000 ppm were attributed as toxic, and ≥ 1000 ppm was not toxic.

Extraction and Isolation

The powdered and dried pod peels of *T.vogelii* (35 g) were extracted to yield a crude methanol extract (18.2 g). These products were then subjected to vacuum liquid chromatography over silica gel 60 GF254, and was eluted with *n*-hexane-EtOAc (10:0–0:10) in a polarity gradient manner (Syah, *et al.*, 2006), and a total of fourteen (I–XIV) fractions were produced. Subsequently,

fraction I was purified through centrifugal chromatography and eluted with chloroform-EtOAc (8:2) to generate isolonchocarpin (80 mg), and the structures were elucidated based on NMR data, and also through comparison with reported spectra values (Athipornchai, *et al.*, 2008).

3. RESULTS AND DISCUSSION

The isolates were obtained as a pure colourless and transparent compounds in the form of needle crystals, with a melting point of 114–115 °C. Table 3 shows the NMR spectrum, featuring the signals of two proton *double doublets* at 2.84 δ_H and 3.00 ppm, respectively assigned to the H3 α and H3 β . Also, one proton signal *double doublets* was observed at δ_H 5.47 ppm for H-2, and two emerged from the two shielding carbons at δ_C 79.8 and 44.4 ppm for (C-2 and C-3). These were further determined as characteristic for flavanone group compounds. The nature of this structure was also confirmed by the presence of a carbon signal C = O ketone conjugated at δ_C 190.6 ppm (Athipornchai, 2008). In addition, the ¹³C NMR spectrum also demonstrated eighteen signals for twenty carbon atoms, including for one ring unit dimethyl tetrahydropyran (δ_C 116.0; 129.0; 77.6; 28.5; 28.2 ppm), and two oxyaryl (δ_C 159.7; 157.7 ppm). Table 3 shows the ¹H NMR spectrum, indicating the presence of an aromatic signal for five protons at δ_H 7.48; 7.43 and 7.39 ppm, corresponding to the unsubstituted B ring. The two oxygen functional groups are further confirmed to occur in ring A, located at C-8a and C-7, according to the prevalence of the oxygen pattern. This phenomenon facilitates the formation of dimethyl tetrahydropyran ring at C-8 or C-6. In addition, the presence of an ortho-coupled aromatic as a *doublet* proton signal ($J = 8.6$ Hz) at δ_H 7.75 and 6.51 ppm respectively denote chemical shifts initiated at C-8. The HMBC spectrum shows the multiplicities and the weak coupling constants between the signal from the aromatic doublet proton at δ_H 7.75 ppm (H-5) and a C = O ketone conjugated carbon at δ_C 190.6 ppm (C-4), Also, a correlated was established between the aromatic double protons at δ_H 6.51 ppm (H-6) with oxyaryl carbon signals at δ_C 159.7 ppm (C-7), alongside two quaternary carbons at δ_C 109.5 and 114.8 ppm, respectively situated by C-8 and C-4a. According to the

coupling constant in H-2 / H-3, the configurations at C-2 and C-3 were determined to be *trans* ($J = 13.2$ Hz), with absolute stereochemistry assumed to follow the prevalence of 2*S*, 3*R*-flavanone. Table 3 shows the evidence of this outcome, indicated by the coupling constant in H-2 / H-3, as (J H-2 / H-3 $\beta = 13.2$ Hz), (J H-2 / H-3 $\alpha = 3.0$ Hz) and (J_{gem} H-3 α / H-3 $\beta = 16.8$ Hz). Based on the spectroscopic data comparison with previous research conducted Athipornchai, *et al.* (2008), the compound is identified as isolonchocarpin.




Figure 2. Chemical structure of isolonchocarpin

Table 3. NMR data for isolonchocarpin

Position	δ_H (mult., J Hz) ppm isolonchocarpin	δ_C (ppm)		HMBC ($^1H \leftrightarrow ^{13}C$)	
		isolonchocarpin	isolonchocarpin*	2J	3J
2	5.47 (<i>dd</i> , 3.0, 13.2)	79.8	79.8	2'	4
3 β	3.00 (<i>dd</i> , 13.2, 16.9)	44.4	44.4	2, 4	1'
3 α	2.84 (<i>dd</i> , 3.0, 16.8)	44.4	44.4	4	4a
4	-	190.6	190.6	-	-
4a	-	114.8	114.8	-	-
5	7.75 (<i>d</i> , 8.6)	127.9	127.9	-	4
6	6.51 (<i>d</i> , 8.6)	111.3	111.3	8, 4a	-
7	-	159.7	159.7	-	-
8	-	109.5	109.5	-	-
8a	-	157.7	157.7	-	-
1'	-	139.1	139.0	-	-
2'	7.48 (<i>dd</i> , 2.0, 7.9)	126.1	126.0	3'	2, 6'
3'	7.43 (<i>dd</i> , 7.9, 8.2)	128.9	128.8	-	5'
4'	7.39 (<i>dt</i> , 2.0, 8.2)	128.7	128.6	3', 5'	-
5'	7.43 (<i>dd</i> , 7.9, 8.2)	128.9	128.8	-	-
6'	7.48 (<i>dd</i> , 2.0, 7.9)	126.1	126.0	5'	2'
4''	6.66 (<i>d</i> , 10.1)	116.0	115.9	8	7, 8a
5''	5.57 (<i>d</i> , 10.1)	129.0	128.9	6''	8
6''	-	77.6	77.6	-	-
7''-CH ₃	1.47 (<i>s</i>)	28.2	28.1	-	5'', 8''
8''-CH ₃	1.45 (<i>s</i>)	28.5	28.4	6''	-

*Athipornchai *et al.* (2008); 1H (300 MHz); ^{13}C (75 MHz), CDCl₃

Table 4. LC₅₀ value of methanol extract and *n*-hexane fraction with maceration method

Extract / Fraction	Replication	LC ₅₀ (ppm)	Mean \pm SD (ppm)
Methanol	1	42.05	41.40 \pm 0.63
	2	41.38	
	3	40.78	
<i>n</i> -Hexane:EtOAc (10:0)	1	37.05	38.16 \pm 1.41
	2	38.41	
	3	36.53	




Figure 3. Selected Heteronuclear multiple-bond correlation for isolonchocarpin Arrows point from carbon to proton

The average LC_{50} value obtained through maceration method was 41.40 ppm, with a deviation standard of 0.63. This result indicates the potential toxic effect of *T. vogelii* pod peels methanol extract at below 1,000 ppm (Meyer *et al.*, 1982). Based on statistical analysis, the *n*-hexane fraction (fraction I) yielded an LC_{50} of 38.16 at 1.41 ppm deviation standard. Therefore, the results indicate the relatively higher toxicity in fraction I compared to the methanol extract, due to the smaller value. Furthermore, the pure isolonchocarpin subsequently isolated from Fraction I is assumed to also demonstrate toxicity.

4. CONCLUSION

This study highlights valuable data on the cytotoxic effect of *T. vogelii* pod peels. The flavanone investigated and identified from the *n*-hexane fraction was isolonchocarpin, which demonstrated an LC_{50} with greater toxicity than the methanol extract. Hence, this yield is assumed to also possess strong cytotoxic property. However, further evaluation is needed to determine further effects, in order to provide the complete safety profile as a phytotherapeutic agent for generally recommended use.

ACKNOWLEDGMENTS

This research was funded by the Research University Grant from Universitas Jenderal Achmad Yani, (Hibah Penelitian Kompetitif Unjani TA.2019/2020, by Valentina Adimurti Kusumaningtyas). The authors are grateful to Dr. Elvira Hermawati,

for the NMR measurements.

REFERENCES

- Achmad SA, Hakim EH, Makmur L, Syah YM, Juliawaty LD, Mujahidin D. 2007. Tumbuhan Obat Indonesia. Bandung(ID): Penerbit ITB, Kebudayaan, Universitas Terbuka: 107-119.
- Andrei CC, Ferreira DT, Faccione M, de Moraes LAB, de Carvalho MG, Braz-Filho R. 2000. C-prenylflavonoids from roots of *Tephrosia tunicata*. *Phytochemistry*. 55: 799-804.
- Andrei CC, Vieira P, Fernandes JB, Da Silva MFDGF, Fo ER. 1997. Dimethylchromene Rotenoids From *Tephrosia candida*. *Phytochemistry*. 6: 1081-1085.
- Athipornchai A, Pancharoen O. 2008. Chemical Constituents From Leaves and Pods of *Millettia brandisiana*. Thesis Master of Science. Department of Chemistry. Silpakorn University, 60.
- Chang LC, Chávez D, Song LL, Farnsworth NR, Pezzuto JM, Kinghorn AD. 2000. Absolute configuration of novel bioactive flavonoids from *Tephrosia purpurea*. *Organic letters*. 2: 515-518.
- Dagne E, Mammo W, Sterner O. 1992. Flavonoids of *Tephrosia polyphylla*. *Phytochemistry*. 31: 3662-3663.
- Dagne E, Bekele A, Waterman PG. 1989. The flavonoids of *Millettia ferruginea* subsp. Darassana in Ethiopia. *Phytochemistry*. 28: 3207-9.
- Ganapaty S, Lakshminarayana K, Lakshmi P, Thomas PS. 2009. Antiprotozoal and cytotoxicity assays of the isolates of *Tephrosia tinctoria*. *Asian Journal of Chemistry*. 21: 1007-1010.
- Gomez-Garibay F, Tellez-Valdez O, Moreno-Torres, Calderon JS. 2002. Flavonoids from *Tephrosia major*. A new Prenyl- β -hydroxychalcone. *Z. Naturforsch.* 57c: 579-583.
- Jang DS, Park JU, Kang YH, Hawthorne ME, Vigo JS, Graham JS, Pezzuto JM, Kinghorn AD. 2003. Potential cancer chemopreventive flavonoids from the stem of *Tephrosia toxicaria*. *J. Nat. Prod.*, 66: 1166-1170.
- Kishore PH, Mopuru VBR, Duvvuru G, Madugula MM, Cristelle C, Bernard B. 2003. A new

- coumestan from *Tephrosia calophylla*. *Chemical Pharmacognosy Bull.*, 51: 194-196.
- Kusumaningtyas VA, Syah YM, Juliawaty LD. 2020. Two stilbenes from Indonesian *Cassia grandis* and their antibacterial activities. *Research Journal of Chemistry and Environment*. 24(1): 61-63.
- Melina, Putra EK, Witanti W, Sukrido, Kusumaningtyas VA. 2020. Design and Implementation of Multi Knowledge Base Expert System Using the SQL Inference Mechanism for Herbal Medicine. *Journal of Physics: Conference Series*. 1477(2): 1-9. <https://doi.org/10.1088/1742-6596/1477/2/022007>
- Meyer BN, Ferrigni NR, Putnam JE, Jacobsen LB, Nichols DE, McLaughlin JL. 1982. Brine shrimp. A convenient general bioassay for active plant constituents. *Planta Med*. 45: 31-4.
- Murthy MSR, Rao VE. 1985. Maxima isoflavone J: a new 0-prenylated isoflavone from *Tephrosia maxima*. *Journal of Natural Products*. 48: 967-968.
- Padmapriya R, Ashwini S, Raveendran R. 2017. In vitro antioxidant and cytotoxic potential of different parts of *Tephrosia purpurea*. *Research in Pharmaceutical Sciences*. 12(1): 31-37.
- Prabhakar P, Vanangamudi A, Gandhidasan R, Venkateswara PR. 1996. Hookerianin: a flavone from *Tephrosia hookeriana*. *Phytochemistry*. 43: 315-316.
- Prashant A, David KGL. 1993. Dehydro-6-hydroxyrotenoid and lupenone from *Tephrosia villosa*. *Phytochemistry*. 32: 484-486.
- Polhill RM, Ravan PH, Stirton CH. 1981. Evolution and Systematics of Evolution of Leguminosae, in *Advances in Legume Systematics*. Part I. Royal Botanic Gardens, Kew, 1.
- Rao EV, Rajendra PM, Sree RM. 1994. A prenylated flavanone from *Tephrosia maxima*. *Phytochemistry*. 37: 111-112.
- Rao EV, Venkataratnam G, Vilain C. 1985. Flavonoids from *Tephrosia fulvinervis*. *Phytochemistry*. 24: 2427-2430.
- Roy M, Mitra SR, Bhattacharyya A, Adityachaudhury N. 1986. Candidone, a flavanone from *Tephrosia candida*. *Phytochemistry*. 25: 961-962.
- Sahgal G, Ramanathan S, Sasidharan S, Mordi MN, Ismail S, Mansor SM. 2010. Brineshrimp lethality and acute oral toxicity studies on *Swietenia mahagoni* (Linn.) Jacq. Seedmethanolic extract. *Pharmacognosy Research*. 2: 25-220.
- Sasidharan S, Darah I, Jain K. 2008. In vivo and In vitro toxicity study of *Gracilaria changii*. *Pharm Biol*. 46: 413-7.
- Syah YM, Hakim EH, Makmur L, Kurdi VA, Ghisalberti EL, Aimi N, Achmad SA. 2006. Prenylated 2-arylbenzofurans from two species of *Artocarpus*. *Natural Product Communications*. 1(7): 549-552. <https://doi.org/10.1177/1934578x0600100706>
- Tanaka T, Iinuma M, Yuki K, Fujii Y, Mizuno M. 1992. Flavonoids in root bark of *Pongamia pinnata*. *Phytochemistry*. 31: 993-998.
- Tarus PK, Machocho AK, Lang'at-Thoruwa CC, Chhabra SC. 2002. Flavonoids from *Tephrosia aequilata*. *Phytochemistry*. 60: 375-379.
- Touqeer S, Saeed MA, Ajaib M. 2013. A review on the phytochemistry and pharmacology of Genus *Tephrosia*. *Phytopharmacology*. 4(3): 598-637.
- W Kalala, A Mwakigonja, S Maregesi, Z. Msengwa, R Mahunnah. 2015. Brine shrimp lethality and acute oral toxicity of *Commiphora swynertonii* (Burrt) exudate. *Pyrex Journal of Medicinal Plant Research*. 1(3): 010-018.
- Yenesew A, Dagne E, Waterman PG. 1989. Flavonoids from the seed pods of *Tephrosia pumila*. *Phytochemistry*. 28: 1291-1292.